

INTRINSIC NOTION OF PRINCIPAL SYMBOL FOR THE HEISENBERG CALCULUS

RAPHAËL PONGE

ABSTRACT. In this paper we define an intrinsic notion of principal for the Hypoelliptic calculus on Heisenberg manifolds. More precisely, the principal symbol of a Ψ_H DO appears as a homogeneous section over the linear dual of the tangent Lie algebra bundle of the manifold. This definition is an important step towards using global K -theoretic tools in the Heisenberg setting, such as those involved in the elliptic setting for proving the Atiyah-Singer index theorem or the regularity of the eta invariant. On the other hand, the intrinsic definition of the principal symbol enables us to give an intrinsic sense to the model operator of Ψ_H DO at point and to give a definitive proof that the Heisenberg calculus is modelled at each point by the calculus of left-invariant Ψ DO's on the tangent group at the point. This also allows us to define an intrinsic Rockland condition for Ψ_H DO's which is shown to be equivalent to the invertibility of the principal symbol, provided that the Levi form has constant rank. Furthermore, we review the main hypoellipticity results on Heisenberg manifolds in terms of the results of the paper. In particular, we complete the treatment of the Kohn Laplacian of [BG] and establish that for the horizontal sublaplacian the invertibility of the principal symbol is equivalent to some condition on the Levi form, called condition $X(k)$. Incidentally, this paper provides us with a relatively up-to-date overview of the main facts about the Heisenberg calculus.

1. INTRODUCTION

The aim of this paper is to define a global and intrinsic notion of the principal symbol for the hypoelliptic calculus on Heisenberg manifolds, along with some applications. On the way, we complete the treatment of the Kohn Laplacian of [BG] and gives a complete treatment for the horizontal sublaplacian on a Heisenberg manifold acting on forms.

Recall that a Heisenberg manifold (M, H) consists of a manifold M together with a distinguished hyperplane bundle $H \subset TM$. This definition includes as main examples the Heisenberg group and its quotients by cocompact lattices, (codimension 1) foliations, CR and contact manifolds, as well as the confoliations of [ET]. Moreover, in this context the natural operators such as Hörmander's sum of squares, the Kohn Laplacian, the horizontal sublaplacian or the contact Laplacian of [Ru], while they may be hypoelliptic, are definitely not elliptic. Thus, the standard elliptic calculus cannot be used efficiently to study these operators.

The relevant substitute to the elliptic calculus is the Heisenberg calculus of Beals-Greiner [BG] and Taylor [Tay], which extends seminal works of Boutet de Monvel [Bo1], Dynin ([Dy1], [Dy2]) and Folland-Stein [FS] (see also [BGH], [CGGP], [EMM], [Gri], [Hö3]). The idea in the Heisenberg calculus, which goes back to Elias Stein, is to build a pseudodifferential calculus on Heisenberg manifolds modelled on that of left-invariant pseudodifferential operators on nilpotent groups. This stems from the fact that the relevant tangent structure for a Heisenberg manifold (M, H) is that of a bundle GM of two-step nilpotent Lie groups (see [BG], [Be], [EMM], [FS], [Gro], [Po2], [Ro2]).

In the original monographs [BG] and [Tay] the principal symbol is only defined in local coordinates, so the definition *a priori* depends on the choice of these coordinates. In the special case of a contact manifold, an intrinsic definition have been given in the unpublished book [EMM], as a section over a bundle of jets of vector fields representing the tangent group bundle of the contact

2000 *Mathematics Subject Classification.* Primary 58J40; Secondary 32V05, 35H10, 53D10.

Key words and phrases. Heisenberg calculus, principal symbol, hypoelliptic operators, CR and contact operators.

manifold. This approach is similar to that of Melrose [Me2] in the setting of the b -calculus for manifolds with boundary.

In this paper we give an intrinsic definition of the principal symbol, valid for an arbitrary Heisenberg manifold, in terms of the description of the tangent Lie group bundle in [Po2]. As a consequence, we can reformulate in a global fashion previously known criteria for existence of parametrices in the Heisenberg calculus. In particular, we can define a Rockland condition for operators in the Heisenberg calculus in a fully intrinsic way and show that this condition allows to determine the existence of a parametrix when the Levi form of the Heisenberg manifold has constant rank.

More importantly, since our approach of the principal symbol connects nicely with the construction of the tangent groupoid of a Heisenberg manifold in [Po2], this presumably allows us to make use of global K -theoretic arguments in the Heisenberg setting, as those involved in the proof of the (full) Atiyah-Singer index theorem ([AS1], [AS2]) and the regularity of the eta invariant for general selfadjoint elliptic Ψ DO's ([APS], [Gi]) or, equivalently, the vanishing of the noncommutative residue of a Ψ DO projection (see [Wo1], [BL], [Po1]). More precisely, this paper can also be seen as a step towards extending the aforementioned results to the Heisenberg setting.

On the other hand, for strictly pseudoconvex CR manifolds this should allow us to recover and to extend in a rather simple way a recent result of Hirachi [Hi] on the invariance of the integral of the logarithmic singularity of the Szegő kernel (see also [Bo2]), in connection with the program of Fefferman [Fe] for the Szegő and the Bergman kernels on strictly pseudoconvex complex domains.

1.1. Intrinsic notion of principal symbol for the Heisenberg calculus. Given a Heisenberg manifold (M, H) and a vector bundle $\mathcal{E}$ we let $\mathfrak{g}^*M$ denote the linear dual of the Lie algebra bundle associated to the tangent Lie group bundle. For $m \in \mathbb{C}$ we let $\Psi_H^m(M, \mathcal{E})$ denote the class of Ψ_H DO's of order m acting on sections of $\mathcal{E}$ and, letting $\pi : \mathfrak{g}^*M \rightarrow M$ be the canonical projection, we define $S_m(M, \mathcal{E})$ as the space of sections $p_m \in C^\infty(\mathfrak{g}^*M \setminus 0, \text{End } \pi_*\mathcal{E})$ such that for any $\lambda > 0$ we have

$$(1.1) \quad p_m(\lambda^2 \xi_0, \lambda \xi') = \lambda^m p_m(\xi_0, \xi'), \quad \xi_0 \in (TM/H)^*, \quad \xi' \in H^*.$$

The principal symbol of an operator $P \in \Psi_H^m(M, \mathcal{E})$ is obtained as an element of $S_m(\mathfrak{g}^*M, \mathcal{E})$ as follows. As shown in [Po2] the tangent Lie group bundle GM of a Heisenberg manifold (M, H) can be described as the bundle $(TM/H) \oplus H$ endowed with a law group encoded by the Levi form $\mathcal{L} : H \times H \rightarrow TM/H$ such that, for sections X and Y of H , we have

$$(1.2) \quad \mathcal{L}(X, Y) = [X, Y] \bmod H.$$

It is also shown in [Po2] that at a point $x \in M$ this approach to GM is equivalent to the extrinsic one in [BG] in terms of the Lie group of a nilpotent Lie algebra of jets of vector fields at point of a local chart. This is made by means of a special kind of privileged coordinates at x , called Heisenberg coordinates.

As a byproduct of the equivalence between the two approaches, we obtain that in Heisenberg coordinates a Heisenberg diffeomorphism $\phi : (M, H) \rightarrow (M', H')$ is well approximated by the induced isomorphism $\phi'_H : GM \rightarrow GM'$ between the tangent Lie group bundles (see [Po2]). This allows us to carry out the proof of the invariance by Heisenberg diffeomorphisms of the Heisenberg calculus in Heisenberg coordinates, rather than in privileged coordinates as in [BG] (see Section 8). As a consequence we can prove:

Theorem 1.1. *Let $P : C_c^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a Ψ_H DO of order m . Then there exists a unique symbol a unique symbol $\sigma_m(P) \in S_m(\mathfrak{g}^*M, \mathcal{E})$ such that, for any $a \in M$, the symbol $\sigma_m(P)(a, \cdot)$ agrees in trivializing Heisenberg coordinates centered at a with the principal symbol of P at $x = 0$.*

We call the symbol $\sigma_m(P)(x, \xi)$ the principal symbol of P . In order to distinguish it from the other definition of the principal symbol in local coordinates, we will sometimes refer to it as the *global* principal symbol, while the other principal symbol will be called the *local* principal symbol.

Let us now describe the main properties of the global principal symbol. First, we have:

Proposition 1.2. *For every $m \in \mathbb{C}$ the principal symbol map $\sigma_m : \Psi_H^m(M, \mathcal{E}) \rightarrow S_m(\mathfrak{g}^*M, \mathcal{E})$ gives rise to a linear isomorphism $\Psi_H^m(M, \mathcal{E})/\Psi_H^{m-1}(M, \mathcal{E}) \xrightarrow{\sim} S_m(\mathfrak{g}^*M, \mathcal{E})$.*

If $P \in \Psi_H^m(M, \mathcal{E})$ has principal symbol $\sigma_m(P)$ then for any $a \in M$ we define the model operator P^a of P at a as the operator as the left-invariant Ψ_H DO on G_aM with symbol $\sigma_m(a, \cdot)$, that is, the left-convolution operator with the inverse Fourier transform of $\sigma_m(a, \cdot)$ (see Definition 4.7).

On the other hand, for any $a \in M$ the product law on G_aM defines a product for symbols,

$$(1.3) \quad *^a : S_{m_1}(\mathfrak{g}_a^*M) \times S_{m_2}(\mathfrak{g}_a^*M) \longrightarrow S_{m_1+m_2}(\mathfrak{g}_a^*M).$$

This product depends smoothly enough on a to give rise to a product

$$(1.4) \quad * : S_{m_1}(\mathfrak{g}^*M, \mathcal{E}) \times S_{m_2}(\mathfrak{g}^*M, \mathcal{E}) \longrightarrow S_{m_1+m_2}(\mathfrak{g}^*M, \mathcal{E}),$$

such that for $p_j \in S_{m_j}(\mathfrak{g}^*M, \mathcal{E})$, $j = 1, 2$, we have

$$(1.5) \quad p_{m_1} * p_{m_2}(a, \xi) = [p_{m_1}(a, \cdot) *^a p_{m_2}(a, \cdot)](\xi) \quad \forall (a, \xi) \in \mathfrak{g}^*M \setminus 0.$$

This comes from the fact that in local coordinates the above product is nicely related to the product of local homogeneous symbols of [BG]. As a consequence we get:

Proposition 1.3. *For $j = 1, 2$ let $P_j \in \Psi_H^{m_j}(M, \mathcal{E})$ and suppose that P_1 or P_2 is properly supported.*

- 1) *We have $\sigma_{m_1+m_2}(P_1 P_2) = \sigma_{m_1}(P) * \sigma_{m_2}(P)$.*
- 2) *At every point $a \in M$ the model operator of $P_1 P_2$ is $(P_1 P_2)^a = P_1^a P_2^a$.*

Next, it is shown in [BG] that the transpose of a Ψ_H DO is again a Ψ_H DO. Thanks to the results of [Po2], we can prove a version of this result in Heisenberg coordinates (see Section 9). As a consequence, we identify the principal symbol of transpose, for we get:

Proposition 1.4. *Let $P \in \Psi_H^m(M, \mathcal{E})$ have principal symbol $\sigma_m(P)$. Then:*

- 1) *The principal symbol of the transpose P^t is $\sigma_m(P^t)(x, \xi) = \sigma_m(x, -\xi)^t$;*
- 2) *If P^a is the model operator of P at a , then the model operator of P^t at a is the transpose operator $(P^a)^t : \mathcal{S}_0(G_x M, \mathcal{E}_x^*) \rightarrow \mathcal{S}_0(G_x M, \mathcal{E}_x^*)$.*

Assume now that M is endowed with a positive density and $\mathcal{E}$ with a Hermitian metric, and let $L^2(M, \mathcal{E})$ be the associated L^2 -Hilbert space. Then we have:

Proposition 1.5. *Let $P \in \Psi_H^m(M, \mathcal{E})$ have principal symbol $\sigma_m(P)$. Then:*

- 1) *The principal symbol of the adjoint P^* is $\sigma_m(P^*)(x, \xi) = \sigma_m(P)(x, \xi)^*$.*
- 2) *If P^a denotes the model operator of P at $a \in M$ then the model operator of P^* at a is the adjoint $(P^a)^*$ of P^a .*

1.2. Rockland condition and hypoellipticity. It is shown in [BG] that, in local coordinates, the invertibility of the local principal symbol is equivalent to the existence of a Ψ_H DO parametrix. Thanks to Proposition 1.3 and the relationship in local coordinates between the local and global principal symbols, we can reformulate this result in a global fashion. More precisely, we have:

Proposition 1.6. *Let $P : C_c^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a Ψ_H DO of order m . Then the following are equivalent:*

- 1) *The principal symbol $\sigma_m(P)$ of P is invertible with respect to the convolution product for homogeneous symbols;*
- 2) *The operator P admits a parametrix Q in $\Psi_H^{-m}(M, \mathcal{E})$.*

Furthermore, if 1) and 2) hold then P is hypoelliptic with gain of $\frac{k}{2}$ -derivatives.

In fact, it may be difficult to determine whether the principal symbol of a Ψ_H DO is invertible because in general the convolution product for symbols is not the pointwise product of symbols. In particular, in the Heisenberg setting, the product of principal symbols is neither commutative, nor microlocal.

Nevertheless, to determine the invertibility of the principal symbol we can make use of a general representation theoretic criterion, the Rockland condition: a Ψ_H DO-operator P satisfies the Rockland condition at a point $a \in M$ when its model operator P^a satisfies the Rockland condition on $G_a M$, i.e., for every nontrivial irreducible unitary representation π of $G_a M$ some (unbounded) operator πP^a acting on the representation space of π is injective on the space of smooth vectors of π .

It is well known that a left-invariant homogeneous Ψ DO on a nilpotent group is hypoelliptic if, and only if, it satisfies the Rockland condition, and it further admits an inverse if, and only if, together with its transpose it satisfies the Rockland condition (e.g. [HN1], [HN2], [CGGP]). Thus if P is a Ψ_H DO such that P and P^t satisfies the Rockland condition at every point then, for any $x \in M$, then the model operator of P^x is invertible. However, whether this inverse depends smoothly enough with respect to x to yield an inverse for the principal symbol is a more delicate issue.

We show here that this occurs when the Levi form (1.2) has constant rank, say $2n$. In this case the tangent Lie group is a fiber bundle with typical fiber $\mathbb{H}^{2n+1} \times \mathbb{R}^{d-2n}$. This allows us to make use of results of Christ *et al.* [CGGP] about families of Ψ DO's on a *fixed* nilpotent graded group to get:

Proposition 1.7. *Let P be a Ψ_H DO of order m , $\Re m \geq 0$, and suppose that the Levi form (1.2) has constant rank. Then the following are equivalent:*

- (i) P and P^t satisfy the Rockland condition at every point of M ;
- (ii) P and P^* satisfy the Rockland condition at every point of M ;
- (iii) The principal symbol of P is invertible.

In particular, if (i) and (ii) holds then P admits a Ψ_H DO parametrix and is hypoelliptic with loss of $\frac{1}{2}\Re m$ derivatives.

In particular, when P is selfadjoint and the Levi form has constant rank the Rockland condition for P is equivalent to the invertibility of the principal symbol of P .

Finally, if $2n$ denotes the rank of the Levi form at a then we have $G_a M \simeq \mathbb{H}^{2n+1} \times \mathbb{R}^{d-2n}$. Therefore, the irreducible unitary representations of $G_a M$ are two kinds, one dimensional representations on $\mathbb{C}$ and infinite dimensional representations on $L^2(\mathbb{R}^n)$. In the former case the Rockland condition corresponds to the invertibility of the principal symbol along H^* , while in the latter case it is enough to look at the representations coming from that of $\mathbb{H}^{2n+1}$ with Planck constants ± 1 (see Section 5).

1.3. Hypoellipticity criterions for sublaplacians. It is interesting to look at the previous results in the case of sublaplacians, as this covers several important examples such as Hörmander's sum of squares, the Kohn Laplacian or the horizontal sublaplacian.

Here by sublaplacian we mean a differential operator $\Delta : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ such that near every point $a \in M$ we can write Δ in the form,

$$(1.6) \quad \Delta = -(X_1^2 + \dots + X_d^2) - i\mu(x)X_0 + O_H(1),$$

where $X_0, X_1, \dots, X_d$ is a H -frame of TM , so that $X_1, \dots, X_d$ span H , the term $\mu(x)$ is a smooth section of $\text{End } \mathcal{E}$ and the notation $O_H(1)$ stands for a differential operator of Heisenberg order ≤ 1 .

In this case the Rockland condition and the invertibility of the reduces to the following. Let $L(x) = (L_{jk}(x))$ be the matrix of $\mathcal{L}$ with respect to the H -frame $X_0, X_1, \dots, X_d$, so that for

$j, k = 1, \dots, d$ we have

$$(1.7) \quad \mathcal{L}(X_j, X_k) = [X_j, X_k] = L_{jk}(x)X_0 \mod H.$$

Let $2n$ be the rank of $\mathcal{L}_a$ and $L(a)$, let $\lambda_1, \dots, \lambda_d$ denoting the eigenvalues of $L(a)$ and consider the condition,

$$(1.8) \quad \text{Sp } \mu(a) \cap \Lambda_a = \emptyset,$$

where the singular set Λ_a is defined as follows,

$$(1.9) \quad \Lambda_a = (-\infty, -\frac{1}{2} \text{Trace } |L(a)|] \cup [\frac{1}{2} \text{Trace } |L(a)|, \infty) \quad \text{if } 2n < d,$$

$$(1.10) \quad \Lambda_a = \{\pm(\frac{1}{2} \text{Trace } |L(a)| + 2\alpha_j |\lambda_j|); \alpha_j \in \mathbb{N}^d\} \quad \text{if } 2n = d.$$

As it turns out this condition makes sense independently of the choice of the H -frame and is the relevant condition to look at in the case of a sublaplacian. More precisely, we have:

Proposition 1.8. *1) The condition (6.23) makes sense intrinsically for any $a \in M$.*

2) At every point $a \in M$ the Rockland conditions for Δ and Δ^t are equivalent to (1.8).

3) The principal symbol of Δ is invertible if, and only if, the condition (1.8) holds at every point of M . Moreover, when the latter occurs Δ admits a parametrix in $\Psi_H^{-2}(M, \mathcal{E})$ and is hypoelliptic with loss of one derivative.

The above result is proved in [BG] in the case of scalar sublaplacians, for which the coefficient $\mu(x)$ in (1.6) is a complex-valued function, but the general case is not dealt with in [BG]. However, it is necessary to extend the result to sublaplacians acting on sections of vector bundles in order to deal with the Kohn Laplacian and the horizontal sublaplacian acting on forms. In particular, Proposition 1.8 allows us to complete the treatment of the Kohn Laplacian in [BG] (see below).

1.4. Main examples of hypoelliptic operators on Heisenberg manifolds. We devote a section where we explain how the Heisenberg calculus, including the results of this paper, provides us with a unified framework to deal with the hypoellipticity of the main geometric operators on a Heisenberg manifold and to recover well-known results proved using various different approaches.

(a) Hörmander's sum of squares. A Hörmander's sum of squares is an operator of the form $\Delta = -(X_1^2 + \dots + X_m^2)$ where $X_1, \dots, X_m$ are vector fields on M . When $X_1, \dots, X_m$ span H we get a sublaplacian and Proposition 1.8 allows us to recover, in this special case, the celebrated result of Hörmander [Hö2] about the hypoellipticity of sum of squares under the bracket condition.

(b) Kohn Laplacian. The Kohn Laplacian is the Laplacian associated to the tangential Cauchy-Riemann complex, or $\bar{\partial}_b$ -complex, on a CR manifold ([KR], [Koh1]). It was shown by Kohn [Koh1] that under the condition $Y(q)$ the Kohn Laplacian acting on (p, q) -forms is hypoelliptic with loss of one derivative.

It was proved by Beals-Greiner [BG] that for the Kohn Laplacian acting on (p, q) -forms the condition (1.8) reduces to the condition $Y(q)$, so we may apply Proposition 1.8 to recover Kohn's result. This allows us to complete the treatment of the Kohn in [BG], because the initial argument there is not quite complete (see Remark 7.3).

(c) Horizontal sublaplacian. The horizontal sublaplacian Δ_b on a Heisenberg manifold (M^{d+1}, H) can be seen as a horizontal Laplacian acting on the horizontal forms, that is on the sections of $\Lambda_{\mathbb{C}}^* H^*$. This operator was first introduced by Tanaka [Ta] for strictly pseudoconvex CR manifolds, but versions of this operator acting on functions were independently defined by Greenleaf [Gr] and Lee [Le].

While Δ_b acting on functions can be seen as a sum of squares up to lower order terms, it seems that little has been done concerning the hypoellipticity of Δ_b acting on forms on a general Heisenberg manifolds, except in the contact case (see [Ta], [Ru]).

We show here that the invertibility of the principal symbol of Δ_b acting on the sections of $\Lambda_{\mathbb{C}}^k H^*$ reduces to a condition involving k and the Levi form of (M, H) only. More precisely, for $a \in M$ let $2n$ be the rank of the Levi form $\mathcal{L}$ at a . We say that the condition $X(k)$ is satisfied at a when

$$(1.11) \quad k \notin \{n, n+1, \dots, d-n\}.$$

We then prove that this condition is the condition (1.8) for Δ_b acting on $\Lambda_{\mathbb{C}}^k H^*$. Thus, using Proposition 1.8 we get:

Proposition 1.9. *Let $\Delta_b : C^\infty(M, \Lambda_{\mathbb{C}}^k H^*) \rightarrow C^\infty(M, \Lambda_{\mathbb{C}}^k H^*)$ be the horizontal sublaplacian acting on horizontal forms of degree k .*

- 1) *At a point $x \in M$ the Rockland condition for Δ_b is equivalent to the condition $X(k)$.*
- 2) *The principal symbol of Δ_b is invertible if, and only if, the condition $X(k)$ is satisfied at every point.*

(d) *Contact Laplacian.* Given a contact manifold (M^{2n+1}, θ) the contact Laplacian is associated to the contact complex defined by Rumin [Ru]. Unlike the previous examples this not a sublaplacian and it is even of order 4 on contact forms of middle degrees.

It has been shown by Rumin [Ru] that the contact Laplacian satisfies the Rockland condition on forms of any degree. Rumin then used results of Helffer-Nourrigat [HN3] to deduce that the contact Laplacian was hypoelliptic maximal. Alternatively, we may use Proposition 1.7 to deduce that on contact forms of any degree the contact Laplacian has an invertible principal symbol, hence admits a parametrix in the Heisenberg calculus and is hypoelliptic.

1.5. Organization of the paper. The paper is organized as follows. In Section 2 we recall the main definitions and examples concerning Heisenberg manifolds and their tangent Lie group bundles. In Section 3 we give a detailed overview of the Heisenberg calculus of [BG] and [Tay], following closely the exposition of [BG].

The section 4 is devoted to the definitions and the main properties of the principal symbols and model operators of Ψ_H DO's. In Section 5 we study the relationships between invertibility of the principal symbol, Rockland condition and hypoellipticity and, in particular, we prove Theorem 1.7.

In Section 6 we deal with sublaplacians and in particular extend the results of [BG] to sublaplacians acting on sections of vector bundles. In Section 7 we deal with the invertibility of the principal symbols of main geometric operators on Heisenberg manifolds: Hörmander's sum of squares, Kohn Laplacian, horizontal sublaplacian and contact Laplacian. In particular, we prove Proposition 1.9.

The last two sections are devoted to the proofs of Proposition 3.18 and Proposition 3.21. These are versions in Heisenberg coordinates of the invariance of the Heisenberg calculus by Heisenberg diffeomorphisms and transposition. The latter were proved in [BG] in a less precise form, but in this paper we need the precise version in Heisenberg coordinates in order to show that the definition of the principal symbol makes sense intrinsically and to determine the principal symbols of the transpose and the adjoint of a Ψ_H DO.

Acknowledgements. The author is grateful to the hospitality of the Mathematics Departments of Princeton University and Harvard University where most part of this paper was written. The research of the author was partially supported by the NSF grant DMS 0409005.

2. HEISENBERG MANIFOLDS AND THEIR TANGENT LIE GROUP BUNDLES

In this section we recall the main facts about Heisenberg manifolds and their tangent Lie group bundle. The exposition here follows closely that of [Po2].

Definition 2.1. 1) A Heisenberg manifold is a smooth manifold M equipped with a distinguished hyperplane bundle $H \subset TM$.

2) A Heisenberg diffeomorphism ϕ from a Heisenberg manifold (M, H) onto another Heisenberg manifold (M', H') is a diffeomorphism $\phi: M \rightarrow M'$ such that $\phi_*H = H'$.

Definition 2.2. Let (M^{d+1}, H) be a Heisenberg manifold. Then:

- 1) A (local) H -frame for TM is a (local) frame $X_0, X_1, \dots, X_d$ of TM so that $X_1, \dots, X_d$ span H .
- 2) A local Heisenberg chart is a local chart together with a local H -frame of TM over its domain.

The main examples of Heisenberg manifolds are the following.

(a) *Heisenberg group.* The $(2n+1)$ -dimensional Heisenberg group $\mathbb{H}^{2n+1}$ is $\mathbb{R}^{2n+1} = \mathbb{R} \times \mathbb{R}^{2n}$ equipped with the group law,

$$(2.1) \quad x.y = (x_0 + y_0 + \sum_{1 \leq j \leq n} (x_{n+j}y_j - x_jy_{n+j}), x_1 + y_1, \dots, x_{2n} + y_{2n}).$$

A left-invariant basis for its Lie algebra $\mathfrak{h}^{2n+1}$ is provided by the vector-fields,

$$(2.2) \quad X_0 = \frac{\partial}{\partial x_0}, \quad X_j = \frac{\partial}{\partial x_j} + x_{n+j} \frac{\partial}{\partial x_0}, \quad X_{n+j} = \frac{\partial}{\partial x_{n+j}} - x_j \frac{\partial}{\partial x_0}, \quad 1 \leq j \leq n,$$

which for $j, k = 1, \dots, n$ and $k \neq j$ satisfy the Heisenberg relations,

$$(2.3) \quad [X_j, X_{n+k}] = -2\delta_{jk}X_0, \quad [X_0, X_j] = [X_j, X_k] = [X_{n+j}, X_{n+k}] = 0.$$

In particular, the subbundle spanned by the vector fields $X_1, \dots, X_{2n}$ defines a left-invariant Heisenberg structure on $\mathbb{H}^{2n+1}$.

(b) *Foliations.* A (smooth) foliation is a manifold M together with a subbundle $\mathcal{F} \subset TM$ which is integrable in the Frobenius' sense, i.e., we have $[\mathcal{F}, \mathcal{F}] \subset \mathcal{F}$. Thus, any codimension 1 foliation is a Heisenberg manifold.

(c) *Contact manifolds.* Opposite to foliations are contact manifolds: a *contact structure* on a manifold M^{2n+1} is given by a global non-vanishing 1-form θ on M such that $d\theta$ is non-degenerate on $H = \ker \theta$. In particular, (M, H) is a Heisenberg manifold. In fact, by Darboux's theorem any contact manifold (M^{2n+1}, θ) is locally contact-diffeomorphic to the Heisenberg group $\mathbb{H}^{2n+1}$ equipped with its standard contact form $\theta^0 = dx_0 + \sum_{j=1}^n (x_j dx_{n+j} - x_{n+j} dx_j)$.

(d) *Confoliations.* According to Elyashberg-Thurston [ET] a *confoliation structure* on an oriented manifold M^{2n+1} is given by a global non-vanishing 1-form θ on M such that $(d\theta)^n \wedge \theta \geq 0$. In particular, when $(d\theta)^n \wedge \theta = 0$ (resp. $(d\theta)^n \wedge \theta > 0$) we are in presence of a foliation (resp. a contact structure). In any case the hyperplane bundle $H = \ker \theta$ defines a Heisenberg structure on M .

(e) *CR manifolds.* A CR structure on an orientable manifold M^{2n+1} is given by a rank n complex subbundle $T_{1,0} \subset T_{\mathbb{C}}M$ which is integrable in Frobenius' sense and such that $T_{1,0} \cap T_{0,1} = \{0\}$, where $T_{0,1} = \overline{T_{1,0}}$. Equivalently, the subbundle $H = \Re(T_{1,0} \otimes T_{0,1})$ has the structure of a complex bundle of (real) dimension $2n$. In particular, the pair (M, H) forms a Heisenberg manifold.

Moreover, since M is orientable and H is orientable by means of its complex structure, the normal bundle TM/H is orientable, hence admits a global nonvanishing section T . Let θ be the global section of T^*M/H^* such that $\theta(T) = 1$ and θ annihilates H . Then Kohn's Levi form is the form L_θ on $T_{1,0}$ such that, for sections Z and $\bar{W}$ of $T_{1,0}$, we have

$$(2.4) \quad L_\theta(Z, \bar{W}) = -id\theta(Z, \bar{W}) = i\theta([Z, \bar{W}]).$$

We say that M is strictly pseudoconvex (resp. nondegenerate, κ -strictly pseudoconvex) when for some choice of θ the Levi form L_θ is everywhere positive definite (resp. is everywhere non-degenerate, has everywhere signature $(n - \kappa, \kappa, 0)$). In particular, when M is nondegenerate the 1-form θ is a contact form on M .

The main example of a CR manifold is that of the (smooth) boundary $M = \partial D$ of a complex domain $D \subset \mathbb{C}^n$. In particular, when D is strongly pseudoconvex (or strongly pseudoconcave) M is strictly pseudoconvex.

2.1. The tangent Lie group bundle. A simple description of the tangent Lie group bundle of a Heisenberg manifold (M^{d+1}, H) can be given as follows.

Lemma 2.3 ([Po2]). *The Lie bracket of vector fields induces 2-form,*

$$(2.5) \quad \mathcal{L} : H \times H \longrightarrow TM/H,$$

such that, for any $a \in M$ and any sections X and Y of H near a , we have

$$(2.6) \quad \mathcal{L}_a(X(a), Y(a)) = [X, Y](a) \mod H_a.$$

Definition 2.4. *The 2-form $\mathcal{L}$ is called the Levi form of (M, H) .*

The Levi form $\mathcal{L}$ allows us to define a bundle $\mathfrak{g}M$ of graded Lie algebras by endowing $(TM/H) \oplus H$ with the smooth fields of Lie Brackets and gradings such that

$$(2.7) \quad [X_0 + X', Y_0 + Y']_a = \mathcal{L}_a(X', Y') \quad \text{and} \quad t.(X_0 + X') = t^2 X_0 + tX', \quad t \in \mathbb{R},$$

for $a \in M$ and X_0, Y_0 in $T_a M/H_a$ and X', Y' in H_a .

Definition 2.5. *The bundle $\mathfrak{g}M$ is called the tangent Lie algebra bundle of M .*

As we can easily check $\mathfrak{g}M$ is a bundle of 2-step nilpotent Lie algebras which contains the normal bundle TM/H in its center. Therefore, its associated graded Lie group bundle GM can be described as follows. As a bundle GM is $(TM/H) \oplus H$ and the exponential map is the identity. In particular, the grading of GM is as in (2.7). Moreover, since $\mathfrak{g}M$ is 2-step nilpotent the Campbell-Hausdorff formula gives

$$(2.8) \quad (\exp X)(\exp Y) = \exp(X + Y + \frac{1}{2}[X, Y]), \quad X, Y \text{ sections of } \mathfrak{g}M.$$

From this we deduce that the product on GM is such that

$$(2.9) \quad (X_0 + X').(Y_0 + X') = X_0 + Y_0 + \frac{1}{2}\mathcal{L}(X', Y') + X' + Y',$$

for sections X_0, Y_0 of TM/H and sections X', Y' of H .

Definition 2.6. *The bundle GM is called the tangent Lie group bundle of M .*

In fact, the fibers of GM are classified by the Levi form $\mathcal{L}$ as follows.

Proposition 2.7 ([Po2]). *1) Let $a \in M$. Then $\mathcal{L}_a$ has rank $2n$ if, and only if, as a graded Lie group $G_a M$ is isomorphic to $\mathbb{H}^{2n+1} \times \mathbb{R}^{d-2n}$.*

2) The Levi form $\mathcal{L}$ has constant rank $2n$ if, and only if, GM is a fiber bundle with typical fiber $\mathbb{H}^{2n+1} \times \mathbb{R}^{d-2n}$.

Now, let $\phi : (M, H) \rightarrow (M', H')$ be a Heisenberg diffeomorphism from (M, H) onto another Heisenberg manifold (M', H') . Since $\phi_* H = H'$ we see that ϕ' induces a smooth vector bundle isomorphism $\bar{\phi} : TM/H \rightarrow TM'/H'$.

Definition 2.8. *We let $\phi'_H : (TM/H) \oplus H \rightarrow (TM'/H') \oplus H'$ denote the vector bundle isomorphism such that*

$$(2.10) \quad \phi'_H(a)(X_0 + X') = \bar{\phi}'(a)X_0 + \phi'(a)X',$$

for any $a \in M$ and any $X_0 \in T_a/H_a$ and $X' \in H_a$.

Proposition 2.9 ([Po2]). *The vector bundle isomorphism ϕ'_H is an isomorphism of graded Lie group bundles from GM onto GM' . In particular, the Lie group bundle isomorphism class of GM depends only on the Heisenberg diffeomorphism class of (M, H) .*

2.2. Heisenberg coordinates and nilpotent approximation of vector fields. It is interesting to relate the intrinsic description of GM above with the more extrinsic description of [BG] (see also [Be], [EMM], [FS], [Gro], [Ro2]) in terms of the Lie group associated to a nilpotent Lie algebra of model vector fields.

First, let $a \in M$ and let us describe $\mathfrak{g}_a M$ as the graded Lie algebra of left-invariant vector fields on $G_a M$ by identifying any $X \in \mathfrak{g}_a M$ with the left-invariant vector fields L_X on $G_a M$ given by

$$(2.11) \quad L_X f(x) = \frac{d}{dt} f[(t \exp X) \cdot x]_{|t=0} = \frac{d}{dt} f[(tX) \cdot x]_{|t=0}, \quad f \in C^\infty(G_a M).$$

This allows us to associate to any vector fields X near a a unique left-invariant vector fields X^a on $G_a M$ such that

$$(2.12) \quad X^a = \begin{cases} L_{X_0(a)} & \text{if } X(a) \notin H_a, \\ L_{X(a)} & \text{otherwise,} \end{cases}$$

where $X_0(a)$ denotes the class of $X(a)$ modulo H_a .

Definition 2.10. *The left-invariant vector fields X^a is called the model vector fields of X at a .*

Let us look at the above construction in terms of a H -frame $X_0, \dots, X_d$ near a , i.e. of a local trivialization of the vector bundle $(TM/H) \oplus H$. For $j, k = 1, \dots, d$ we let

$$(2.13) \quad \mathcal{L}(X_j, X_k) = [X_j, X_k]X_0 = L_{jk}X_0 \mod H.$$

With respect to the coordinate system $(x_0, \dots, x_d) \rightarrow x_0 X_0(a) + \dots + x_d X_d(a)$ we can write the product law of $G_a M$ as

$$(2.14) \quad x \cdot y = (x_0 + \frac{1}{2} \sum_{j,k=1}^d L_{jk} x_j x_k, x_1, \dots, x_d).$$

Then the vector fields X_j^a , $j = 0, \dots, d$, in (2.12) are the left-invariant vector fields corresponding to the vectors e_j , $j = 0, \dots, d$, of the canonical basis of $\mathbb{R}^{d+1}$, i.e., we have

$$(2.15) \quad X_0^a = \frac{\partial}{\partial x_0} \quad \text{and} \quad X_j^a = \frac{\partial}{\partial x_j} - \frac{1}{2} \sum_{k=1}^d L_{jk} x_k \frac{\partial}{\partial x_0}, \quad 1 \leq j \leq d.$$

In particular, for $j, k = 1, \dots, d$ we have the relations,

$$(2.16) \quad [X_j^a, X_k^a] = L_{jk}(a) X_0^a, \quad [X_j^a, X_0^a] = 0.$$

Now, let $\kappa : \text{dom } \kappa \rightarrow U$ be a Heisenberg chart near $a = \kappa^{-1}(u)$ and let $X_0, \dots, X_d$ be the associated H -frame of TU . Then there exists a unique affine coordinate change $x \rightarrow \psi_u(x)$ such that $\psi_u(u) = 0$ and $\psi_{u*} X_j(0) = \frac{\partial}{\partial x_j}$ for $j = 0, 1, \dots, d$. Indeed, if for $j = 1, \dots, d$ we set $X_j(x) = \sum_{k=0}^d B_{jk}(x) \frac{\partial}{\partial x_k}$ then we have

$$(2.17) \quad \psi_u(x) = A(u)(x - u), \quad A(u) = (B(u)^t)^{-1}.$$

Definition 2.11. 1) *The coordinates provided by ψ_u are called the privileged coordinates at u with respect to the H -frame $X_0, \dots, X_d$.*

2) *The map ψ_u is called the privileged-coordinate map with respect to the H -frame $X_0, \dots, X_d$.*

Remark 2.12. The privileged coordinates at u are called u -coordinates in [BG], but they correspond to the privileged coordinates of [Be] and [Gro] in the special case of a Heisenberg manifold.

Next, on $\mathbb{R}^{d+1}$ we consider the dilations

$$(2.18) \quad \delta_t(x) = t \cdot x = (t^2 x_0, t x_1, \dots, t x_d), \quad t \in \mathbb{R},$$

with respect to which $\frac{\partial}{\partial x_0}$ is homogeneous of degree -2 and $\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_d}$ is homogeneous of degree -1 .

Since in the privileged coordinates at u we have $X_j(0) = \frac{\partial}{\partial x_j}$ we can write

$$(2.19) \quad X_j = \frac{\partial}{\partial x_j} + \sum_{k=0}^d a_{jk}(x) \frac{\partial}{\partial x_k}, \quad j = 0, 1, \dots, d,$$

where the a_{jk} 's are smooth functions such that $a_{jk}(0) = 0$. Therefore, we may define

$$(2.20) \quad X_0^{(u)} = \lim_{t \rightarrow 0} t^2 \delta_t^* X_0 = \frac{\partial}{\partial x_0},$$

$$(2.21) \quad X_j^{(u)} = \lim_{t \rightarrow 0} t^{-1} \delta_t^* X_j = \frac{\partial}{\partial x_j} + \sum_{k=1}^d b_{jk} x_k \frac{\partial}{\partial x_0}, \quad j = 1, \dots, d,$$

where for $j, k = 1, \dots, d$ we have set $b_{jk} = \partial_{x_k} a_{j0}(0)$.

Observe that $X_0^{(u)}$ is homogeneous of degree -2 and $X_1^{(u)}, \dots, X_d^{(u)}$ are homogeneous of degree -1 . Moreover, for $j, k = 1, \dots, d$ we have

$$(2.22) \quad [X_j^{(u)}, X_0^{(u)}] = 0 \quad \text{and} \quad [X_j^{(u)}, X_0^{(u)}] = (b_{kj} - b_{jk}) X_0^{(u)}.$$

Thus, the linear space spanned by $X_0^{(u)}, X_1^{(u)}, \dots, X_d^{(u)}$ is a graded 2-step nilpotent Lie algebra $\mathfrak{g}^{(u)}$. In particular, $\mathfrak{g}^{(u)}$ is the Lie algebra of left-invariant vector fields over the graded Lie group $G^{(u)}$ consisting of $\mathbb{R}^{d+1}$ equipped with the grading (2.18) and the group law,

$$(2.23) \quad x.y = (x_0 + \sum_{j,k=1}^d b_{kj} x_j x_k, x_1, \dots, x_d).$$

Now, if near a we let $\mathcal{L}(X_j, X_k) = [X_j, X_k] = L_{jk}(x) X_0 \bmod H$, then we get

$$(2.24) \quad [X_j^{(u)}, X_k^{(u)}] = \lim_{t \rightarrow 0} [t \delta_t^* X_j, t \delta_t^* X_k] = \lim_{t \rightarrow 0} t^2 \delta_t^* (L_{jk}(\circ \kappa^{-1}(x)) X_0) = L_{jk}(a) X_0^{(u)}.$$

Comparing this with (2.16) and (2.22) then shows that $\mathfrak{g}^{(u)}$ has the same the constant structures as those of $\mathfrak{g}_a M$, hence is isomorphic to it. Consequently, the Lie groups $G^{(u)}$ and $G_a M$ are isomorphic. In fact, as shown in [BG] and [Po2], an explicit isomorphism is given by

$$(2.25) \quad \phi_u(x_0, \dots, x_d) = (x_0 - \frac{1}{4} \sum_{j,k=1}^d (b_{jk} + b_{kj}) x_j x_k, x_1, \dots, x_d).$$

Definition 2.13. Let $\varepsilon_u = \phi_u \circ \psi_u$. Then:

1) The new coordinates provided by ε_u are called *Heisenberg coordinates at u with respect to the H -frame $X_0, \dots, X_d$* .

2) The map ε_u is called the *u -Heisenberg coordinate map*.

Remark 2.14. The Heisenberg coordinates at u have been also considered in [BG] as a technical tool for inverting the principal symbol of a hypoelliptic sublaplacian.

Next, by [Po2, Lem. 1.17] we have

$$(2.26) \quad \phi_* X_0^{(u)} = \frac{\partial}{\partial x_0} = X_0^a \quad \text{and} \quad \phi_* X_j^{(u)} = \frac{\partial}{\partial x_j} - \frac{1}{2} \sum_{k=1}^d L_{jk} x_k \frac{\partial}{\partial x_0} = X_j^a, \quad j = 1, \dots, d.$$

Since ϕ_u commutes with the Heisenberg dilations (2.18), using (2.20)–(2.21) we get

$$(2.27) \quad \lim_{t \rightarrow 0} t^2 \delta_t^* \phi_{u*} X_0^{(u)} = X_0^a \quad \text{and} \quad \lim_{t \rightarrow 0} t \delta_t^* \phi_{u*} X_j^{(u)} = X_j^a, \quad j = 1, \dots, d.$$

In fact, as shown in [Po2], in Heisenberg coordinates at a for any vector fields X as $t \rightarrow 0$ we have

$$(2.28) \quad \delta_t^* X = \begin{cases} t^{-2} X^a + O(t^{-1}) & \text{if } X(a) \in H_a, \\ t^{-1} X^a + O(1) & \text{otherwise.} \end{cases}$$

Therefore, we obtain:

Proposition 2.15 ([Po2]). *In the Heisenberg coordinates centered at $m = \kappa^{-1}(u)$ the tangent Lie group $G_a M$ coincides with $G^{(u)}$ and for any vector fields X the model vector fields X^a approximates X near a in the sense of (2.28).*

One consequence of the equivalence between the two approaches to GM is a tangent approximation for Heisenberg diffeomorphisms as follows.

Let $\phi : (M, H) \rightarrow (M', H')$ be a Heisenberg diffeomorphism from (M, H) to another Heisenberg manifold (M', H') . We also endow $\mathbb{R}^{d+1}$ with the pseudo-norm,

$$(2.29) \quad \|x\| = (x_0^2 + (x_1^2 + \dots + x_d^2)^2)^{1/4}, \quad x \in \mathbb{R}^{d+1},$$

so that, for any $x \in \mathbb{R}^{d+1}$ and any $t \in \mathbb{R}$, we have

$$(2.30) \quad \|t.x\| = |t| \|x\|.$$

Proposition 2.16 ([Po2, Prop. 2.21]). *Let $a \in M$ and set $m' = \phi(a)$. Then, in Heisenberg coordinates at a and at a' the diffeomorphism $\phi(x)$ has a behavior near $x = 0$ of the form*

$$(2.31) \quad \phi(x) = \phi'_H(0)x + (O(\|x\|^3), O(\|x\|^2), \dots, O(\|x\|^2)).$$

In particular, there are no terms of the form $x_j x_k$, $1 \leq j, k \leq d$, in the Taylor expansion of $\phi_0(x)$ at $x = 0$.

Remark 2.17. An asymptotics similar to (2.31) is given in [Be, Prop. 5.20] in privileged coordinates at u and $u' = \kappa_1(a')$, but the leading term there is only a Lie algebra isomorphism from $\mathfrak{g}^{(u)}$ onto $\mathfrak{g}^{(u')}$. This is only in Heisenberg coordinates that we recover the Lie group isomorphism $\phi'_H(a)$ as the leading term of the asymptotics.

3. HYPOELLIPTIC CALCULUS ON HEISENBERG MANIFOLDS

The Heisenberg calculus is the relevant pseudodifferential tool to study hypoelliptic operators on Heisenberg manifolds. It was independently invented by Beals-Greiner [BG] and Taylor [Tay], extending previous works of Boutet de Monvel [Bo1], Folland-Stein [FS] and Dynin ([Dy1], [Dy2]) (see also [BGH], [CGGP], [EMM], [Gri], [Hö3], [RSt]).

The idea in the Heisenberg calculus is to have a pseudodifferential calculus on a Heisenberg manifold (M, H) which is modeled at any point $a \in M$ by the calculus of left-invariant pseudodifferential operators on the tangent group $G_a M$.

3.1. Left-invariant pseudodifferential operators. Let (M^{d+1}, H) be a Heisenberg manifold and let $G = G_a M$ be the tangent Lie group of M at a point $a \in M$. We recall here the main facts about left-invariant pseudodifferential operators on G (see also [BG], [CGGP], [Tay]).

Recall that for any finite dimensional vector space E the Schwartz class $\mathcal{S}(E)$ is a Fréchet space and the Fourier transform is the continuous isomorphism of $\mathcal{S}(E)$ onto $\mathcal{S}(E^*)$ given by

$$(3.1) \quad \hat{f}(\xi) = \int_E e^{i\langle \xi, x \rangle} f(x) dx, \quad f \in \mathcal{S}(E), \quad \xi \in E^*,$$

where dx denotes the Lebesgue measure of E .

Definition 3.1. $\mathcal{S}_0(E)$ is the closed subspace of $\mathcal{S}(E)$ consisting of $f \in \mathcal{S}(E)$ such that for any differential operator P on E^* we have $(P\hat{f})(0) = 0$.

Since G has the same underlying set as that of its Lie algebra $\mathfrak{g} = \mathfrak{g}_x M$ we can let $\mathcal{S}(G)$ and $\mathcal{S}_0(G)$ denote the Fréchet spaces $\mathcal{S}(E)$ and $\mathcal{S}_0(E)$ associated to the underlying linear space E of $\mathfrak{g}$ (notice that the Lebesgue measure of E coincides with the Haar measure of G since G is nilpotent).

Next, for $\lambda \in \mathbb{R}$ and $\xi = \xi_0 + \xi'$ in $\mathfrak{g}^* = (T_a^* M / H_a^*) \oplus H_a$ we let

$$(3.2) \quad \lambda \cdot \xi = \lambda \cdot (\xi_0 + \xi') = \lambda^2 \xi_0 + \lambda \xi'.$$

Definition 3.2. $S_m(\mathfrak{g}^*)$, $m \in \mathbb{C}$, is the space of functions $p \in C^\infty(\mathfrak{g}^* \setminus 0)$ which are homogeneous of degree m , in the sense that, for any $\lambda > 0$ we have

$$(3.3) \quad p(\lambda \cdot \xi) = \lambda^m p(\xi) \quad \forall \xi \in \mathfrak{g}^* \setminus 0.$$

In addition $S_m(\mathfrak{g}^*)$ is endowed with the Fréchet space topology induced from that of $C^\infty(\mathfrak{g}^* \setminus 0)$.

Note that the image $\hat{\mathcal{S}}_0(G)$ of $\mathcal{S}(G)$ under the Fourier transform consists of functions $v \in \mathcal{S}(\mathfrak{g}^*)$ such that, given any norm $|\cdot|$ on G , near $\xi = 0$ we have $|g(\xi)| = O(|\xi|^N)$ for any integer $N \geq 0$. Thus, any $p \in S_m(\mathfrak{g}^*)$ defines an element of $\hat{\mathcal{S}}_0(\mathfrak{g}^*)'$ by letting

$$(3.4) \quad \langle p, g \rangle = \int_{\mathfrak{g}^*} p(\xi) g(\xi) d\xi, \quad g \in \hat{\mathcal{S}}_0(\mathfrak{g}^*).$$

This allows us to define the inverse Fourier transform of p as the element $\check{p} \in \mathcal{S}_0(G)'$ such that

$$(3.5) \quad \langle \check{p}, f \rangle = \langle p, \check{f} \rangle \quad \forall f \in \mathcal{S}_0(G).$$

Proposition 3.3 ([BG], [CGGP]). 1) For any $p \in S_m(\mathfrak{g}^*)$ the left-convolution by $\check{p}$,

$$(3.6) \quad \check{p} * f(x) := \langle \check{p}(y), f(x \cdot y^{-1}) \rangle, \quad f \in \mathcal{S}_0(G),$$

defines a continuous endomorphism of $\mathcal{S}_0(G)$.

2) There is a continuous bilinear product,

$$(3.7) \quad * : S_{m_1}(\mathfrak{g}^*) \times S_{m_2}(\mathfrak{g}^*) \longrightarrow S_{m_1+m_2}(\mathfrak{g}^*),$$

such that, for any $p_1 \in S_{m_1}(\mathfrak{g}^*)$ and $p_2 \in S_{m_2}(\mathfrak{g}^*)$, the composition of the left-convolution operators by $\check{p}_1$ and $\check{p}_2$ is the left-convolution operator by $(p_1 * p_2)^\vee$, that is, we have

$$(3.8) \quad \check{p}_1 * (\check{p}_2 * f) = (p_1 * p_2)^\vee * f \quad \forall f \in \mathcal{S}_0(G).$$

Let us also mention that if $p \in S_m(\mathfrak{g}^*)$ then the convolution operator $Pu = \check{p} * f$ is a pseudodifferential operator. Indeed, let $X_0(a), \dots, X_d(a)$ be a (linear) basis of $\mathfrak{g}$ so that $X_0(a)$ is in $T_a M / H_a$ and $X_1(a), \dots, X_d(a)$ span H_a . For $j = 0, \dots, d$ let X_j^a be the left-invariant vector fields on G such that $X_j^w|_{x=0} = X_j(a)$. The basis $X_0(a), \dots, X_d(a)$ yields a linear isomorphism $\mathfrak{g} \simeq \mathbb{R}^{d+1}$, hence a global chart of G . In this chart p is a homogeneous symbol on $\mathbb{R}^{d+1} \setminus 0$ with respect to the dilations

$$(3.9) \quad \lambda \cdot x = (\lambda^2 x_0, \lambda x_1, \dots, \lambda x_d), \quad x \in \mathbb{R}^{d+1}, \quad \lambda > 0.$$

Similarly, each vector fields $\frac{1}{i} X_j^a$, $j = 0, \dots, d$, corresponds to a vector fields on $\mathbb{R}^{d+1}$ whose symbol is denoted $\sigma_j^a(x, \xi)$. Then, setting $\sigma = (\sigma_0, \dots, \sigma_d)$, it can be shown that in the above chart the operator P is given by

$$(3.10) \quad Pf(x) = \int_{\mathbb{R}^{d+1}} e^{ix \cdot \xi} p(\sigma^a(x, \xi)) \hat{f}(\xi), \quad f \in \mathcal{S}_0(\mathbb{R}^{d+1}).$$

In other words P is the pseudodifferential operator $p(-iX^a) := p(\sigma^a(x, D))$ acting on $\mathcal{S}_0(\mathbb{R}^{d+1})$.

3.2. Ψ_H DO's on an open subset of $\mathbb{R}^{d+1}$. Let U be an open subset of $\mathbb{R}^{d+1}$ together with a hyperplane bundle $H \subset TU$ and a H -frame $X_0, X_1, \dots, X_d$ of TU . Then the class of Ψ_H DO's on U is a class of pseudodifferential operators modelled on that of homogeneous convolution operators on the fibers of GU .

Definition 3.4. $S_m(U \times \mathbb{R}^{d+1})$, $m \in \mathbb{C}$, is the space of symbols $p(x, \xi) \in C^\infty(U \times \mathbb{R}^{d+1} \setminus 0)$ that are homogeneous of degree m with respect to the ξ -variable, that is,

$$(3.11) \quad p(x, \lambda \xi) = \lambda^m p(x, \xi) \quad \text{for any } \lambda > 0,$$

where $\xi \rightarrow \lambda \xi$ denotes the Heisenberg dilation (3.9).

Observe that the homogeneity of $p \in S_m(U \times \mathbb{R}^{d+1})$ implies that, for any compact $K \subset U$, it satisfies the estimates

$$(3.12) \quad |\partial_x^\alpha \partial_\xi^\beta p(x, \xi)| \leq C_{K\alpha\beta} \|\xi\|^{\Re m - \langle \beta \rangle}, \quad x \in K, \quad \xi \neq 0,$$

where $\|\xi\| = (|\xi_0|^2 + |\xi_1|^4 + \dots + |\xi_d|^4)^{1/4}$ and $\langle \alpha \rangle = 2\alpha_0 + \alpha_1 + \dots + \alpha_d$.

Definition 3.5. $S^m(U \times \mathbb{R}^{d+1})$, $m \in \mathbb{C}$, consists of symbols $p \in C^\infty(U \times \mathbb{R}^{d+1})$ with an asymptotic expansion $p \sim \sum_{j \geq 0} p_{m-j}$, $p_k \in S_k(U \times \mathbb{R}^{d+1})$, in the sense that, for any integer N and for any compact $K \subset U$, we have

$$(3.13) \quad |\partial_x^\alpha \partial_\xi^\beta (p - \sum_{j < N} p_{m-j})(x, \xi)| \leq C_{\alpha\beta NK} \|\xi\|^{\Re m - \langle \beta \rangle - N}, \quad x \in K, \quad \|\xi\| \geq 1.$$

Next, for $j = 0, \dots, d$ let $\sigma_j(x, \xi)$ denote the symbol of $\frac{1}{i}X_j$ (in the classical sense) and set $\sigma = (\sigma_0, \dots, \sigma_d)$. For any $p \in S^m(U \times \mathbb{R}^{d+1})$ it can be shown that the symbol $p_\sigma(x, \xi) := p(x, \sigma(x, \xi))$ is in the Hörmander class of symbols of type $(\frac{1}{2}, \frac{1}{2})$ (see [BG, Prop. 10.22]). Therefore, we define a continuous linear operator from $C_c^\infty(U)$ to $C^\infty(U)$ by letting

$$(3.14) \quad p(x, -iX)f(x) = (2\pi)^{-(d+1)} \int e^{ix \cdot \xi} p(x, \sigma(x, \xi)) \hat{f}(\xi) d\xi, \quad f \in C_c^\infty(U).$$

In the sequel we let $\Psi^{-\infty}(U)$ denotes the class of smoothing operators, i.e. of operators given by smooth kernels.

Definition 3.6. $\Psi_H^m(U)$, $m \in \mathbb{C}$, consists of operators $P : C_c^\infty(U) \rightarrow C^\infty(U)$ of the form

$$(3.15) \quad P = p(x, -iX) + R,$$

with p in $S^m(U \times \mathbb{R}^{d+1})$, called the symbol of P , and with R in $\Psi^{-\infty}(U)$.

The above definition of the symbol of P differs from that of [BG], since there the authors defined it to be $p_\sigma(x, \xi) = p(x, \sigma(x, \xi))$. Note also that p is unique modulo $S^{-\infty}(U \times \mathbb{R}^{d+1})$.

Lemma 3.7. For $j = 0, 1, \dots$ let $p_{m-j} \in S_{m-j}(U \times \mathbb{R}^{d+1})$. Then there exists $P \in \Psi_H^m(U)$ with symbol $p \sim \sum_{j \geq 0} p_{m-j}$. Moreover, the operator P is unique modulo smoothing operators.

The class $\Psi_H^m(U)$ does not depend on the choice of the H -frame $X_0, \dots, X_d$ (see [BG, Prop. 10.46]). Moreover, since it is contained in the class of Ψ DO's of type $(\frac{1}{2}, \frac{1}{2})$ we get:

Proposition 3.8. Let P be a Ψ_H DO of order m on U .

1) P extends to a continuous linear mapping from $\mathcal{E}'(U)$ to $\mathcal{D}'(U)$ and has a distribution kernel which is smooth off the diagonal.

3) Let $k = \Re m$ if $\Re m \geq 0$ and $k = \frac{1}{2}\Re m$ otherwise. Then for any $s \in \mathbb{R}$ the operator $P \in \Psi_H^m(U)$ extends to a continuous mapping from $L_{s, \text{comp}}^2(U)$ to $L_{s-k, \text{loc}}^2(U)$.

3.3. Composition of Ψ_H DO's. Recall that there is no symbolic calculus for Ψ DO's of type $(\frac{1}{2}, \frac{1}{2})$ since the product of two such Ψ DO's needs not be again a Ψ DO of type $(\frac{1}{2}, \frac{1}{2})$. However, the fact that the Ψ_H DO's are modelled on left-invariant pseudodifferential operators allows us to construct a symbolic calculus for Ψ_H DO's.

First, for $j = 0, \dots, d$ let $X_j^{(x)}$ be the leading homogeneous part of X_j in privileged coordinates centered at x defined according to (2.20)–(2.21). These vectors span a nilpotent Lie algebra of left-invariant vector fields on a nilpotent graded Lie group G^x which corresponds to $G_x U$ by pulling back the latter from the Heisenberg coordinates at x to the privileged coordinates at x .

As alluded to above the product law of $G^{(x)}$ defines a convolution product for symbols,

$$(3.16) \quad *^{(x)} : S_{m_1}(\mathbb{R}^{d+1}) \times S_{m_2}(\mathbb{R}^{d+1}) \longrightarrow S_{m_1+m_2}(\mathbb{R}^{d+1}).$$

such that, with the notations of (3.10), on $\mathcal{L}(S_0(\mathbb{R}^{d+1}))$ we have

$$(3.17) \quad p_1(-iX^{(x)})p_2(-iX^{(x)}) = (p_1 *^{(x)} p_2)(-iX^{(x)}) \quad \forall p_j \in S_{m_j}(\mathbb{R}^{d+1}).$$

As it turns out the product $*^{(x)}$ depends smoothly on x (see [BG, Prop. 13.33]). Therefore, we get a continuous bilinear product,

$$(3.18) \quad * : S_{m_1}(U \times \mathbb{R}^{d+1}) \times S_{m_2}(U \times \mathbb{R}^{d+1}) \rightarrow S_{m_1+m_2}(U \times \mathbb{R}^{d+1}),$$

$$(3.19) \quad p_1 * p_2(x, \xi) = (p_1(x, \cdot) *^{(x)} p_2(x, \cdot))(\xi), \quad p_j \in S_{m_j}(U \times \mathbb{R}^{d+1}).$$

Proposition 3.9 ([BG, Thm. 14.7]). *For $j = 1, 2$ let $P_j \in \Psi_H^{m_j}(U)$ have symbol $p_j \sim \sum_{k \geq 0} p_{j, m_j - k}$ and assume that one of these operators is properly supported. Then the operator $P = P_1 P_2$ is a Ψ_H DO of order $m_1 + m_2$ and has symbol $p \sim \sum_{k \geq 0} p_{m_1+m_2-k}$, with*

$$(3.20) \quad p_{m_1+m_2-k}(x, \xi) = \sum_{k_1+k_2 \leq k} \sum_{\alpha, \beta, \gamma, \delta}^{(k-k_1-k_2)} h_{\alpha\beta\gamma\delta}(x) (D_\xi^\delta p_{1, m_1-k_1}) * (\xi^\gamma \partial_x^\alpha \partial_\xi^\beta p_{2, m_2-k_2})(x, \xi),$$

where $\sum_{\alpha\beta\gamma\delta}^{(l)}$ denotes the sum over the indices such that $|\beta| = |\gamma|$ and $|\alpha| + |\beta| \leq \langle \beta \rangle - \langle \gamma \rangle + \langle \delta \rangle = l$, and the functions $h_{\alpha\beta\gamma\delta}(x)$'s are polynomials in the derivatives of the coefficients of the vector fields $X_0, \dots, X_d$.

3.4. The distribution kernels of Ψ_H DO's. An important fact about Ψ DO's is their characterization in terms of their distribution kernels.

First, we extend the notion of homogeneity of functions to distributions. For $K \in \mathcal{S}'(\mathbb{R}^{d+1})$ and for $\lambda > 0$ we let K_λ denote the element of $\mathcal{S}'(\mathbb{R}^{d+1})$ such that

$$(3.21) \quad \langle K_\lambda, f \rangle = \lambda^{-(d+2)} \langle K(x), f(\lambda^{-1} \cdot x) \rangle \quad \forall f \in \mathcal{S}(\mathbb{R}^{d+1}).$$

In the sequel we will also use the notation $K(\lambda \cdot x)$ for denoting $K_\lambda(x)$. We then say that K is homogeneous of degree m , $m \in \mathbb{C}$, when $K_\lambda = \lambda^m K$ for any $\lambda > 0$.

Definition 3.10. $\mathcal{S}'_{reg}(\mathbb{R}^{d+1})$ consists of tempered distributions on $\mathbb{R}^{d+1}$ which are smooth outside the origin. We equip it with the weakest topology such that the inclusions of $\mathcal{S}'_{reg}(\mathbb{R}^{d+1})$ into $\mathcal{S}'(\mathbb{R}^{d+1})$ and $C^\infty(\mathbb{R}^{d+1} \setminus 0)$ are continuous.

Definition 3.11. $\mathcal{K}_m(U \times \mathbb{R}^{d+1})$, $m \in \mathbb{C}$, consists of distributions $K(x, y)$ in $C^\infty(U) \hat{\otimes} \mathcal{S}'_{reg}(\mathbb{R}^{d+1})$ such that for some functions $c_\alpha(x) \in C^\infty(U)$, $\langle \alpha \rangle = m$, we have

$$(3.22) \quad K(x, \lambda \cdot y) = \lambda^m K(x, y) + \lambda^m \log \lambda \sum_{\langle \alpha \rangle = m} c_\alpha(x) y^\alpha \quad \text{for any } \lambda > 0.$$

The interest of considering the distribution class $\mathcal{K}_m(U \times \mathbb{R}^{d+1})$ stems from:

Lemma 3.12 ([BG, Prop. 15.24], [CM, Lem. I.4]). 1) Any $p \in S_m(U \times \mathbb{R}^{d+1})$ agrees on $U \times (\mathbb{R}^{d+1} \setminus 0)$ with a distribution $\tau(x, \xi) \in C^\infty(U) \hat{\otimes} \mathcal{S}'(\mathbb{R}^{d+1})$ such that $\check{\tau}_{\xi \rightarrow y}$ is in $\mathcal{K}_{\hat{m}}(U \times \mathbb{R}^{d+1})$, $\hat{m} = -(m+d+2)$.

2) If $K(x, y)$ is in $\mathcal{K}_{\hat{m}}(U \times \mathbb{R}^{d+1})$ then the restriction of $\hat{K}_{y \rightarrow \xi}(x, \xi)$ to $U \times (\mathbb{R}^{d+1} \setminus 0)$ belongs to $S_m(U \times \mathbb{R}^{d+1})$.

This result is a consequence of the solution to the problem of extending a homogeneous function $p \in C^\infty(\mathbb{R}^{d+1} \setminus 0)$ into a homogeneous distribution on $\mathbb{R}^{d+1}$ and of the fact that for $\tau \in \mathcal{S}'(\mathbb{R}^{d+1})$ we have

$$(3.23) \quad (\hat{\tau})_\lambda = |\lambda|^{-(d+2)} (\tau_{\lambda^{-1}})^\wedge \quad \forall \lambda \in \mathbb{R} \setminus 0.$$

In particular, if τ is homogeneous of degree m then $\hat{\tau}$ is homogeneous of degree $-(m+d+2)$.

The relevant class of kernels for the Heisenberg calculus is the following.

Definition 3.13. $\mathcal{K}^m(U \times \mathbb{R}^{d+1})$, $m \in \mathbb{C}$, consists of distributions $K \in \mathcal{D}'(U \times \mathbb{R}^{d+1})$ with an asymptotic expansion $K \sim \sum_{j \geq 0} K_{m+j}$, $K_l \in \mathcal{K}_l(U \times \mathbb{R}^{d+1})$, in the sense that, for any integer N , as soon as J is large enough we have

$$(3.24) \quad K - \sum_{j \leq J} K_{m+j} \in C^N(U \times \mathbb{R}^{d+1}).$$

Since under the Fourier transform the asymptotic expansion (3.13) for symbols corresponds to that for distributions in (3.24), using Lemma 3.12 we get:

Lemma 3.14 ([BG, pp. 133–134]). Let $K \in \mathcal{D}'(U \times \mathbb{R}^{d+1})$. Then the following are equivalent:

- (i) The distribution K belongs to $\mathcal{K}^m(U \times \mathbb{R}^{d+1})$;
- (ii) We can put K into the form

$$(3.25) \quad K(x, y) = \check{p}_{\xi \rightarrow y}(x, y) + R(x, y),$$

for some $p \in S^{\hat{m}}(U \times \mathbb{R}^{d+1})$, $\hat{m} = -(m+d+2)$, and some $R \in C^\infty(U \times \mathbb{R}^{d+1})$.

Moreover, if (i) and (ii) holds and we expand $K \sim \sum_{j \geq 0} K_{m+j}$, $K_l \in \mathcal{K}_l(U \times \mathbb{R}^{d+1})$, then we have $p \sim \sum_{j \geq 0} p_{\hat{m}-j}$ where $p_{\hat{m}-j} \in S_{\hat{m}-j}(U \times \mathbb{R}^{d+1})$ is the restriction to $U \times (\mathbb{R}^{d+1} \setminus 0)$ of $(K_{m+j})_{y \rightarrow \xi}^\wedge$.

Next, for $x \in U$ let ψ_x denote the affine change to the privileged coordinates at x and let us write $(A_x^t)^{-1}\xi = \sigma(x, \xi)$ with $A_x \in \text{GL}_{d+1}(\mathbb{R})$. Since $\psi_x(x) = 0$ and $\psi_{x*}X_j = \frac{\partial}{\partial y_j}$ at $y = 0$ for $j = 0, \dots, d$, one checks that $\psi_x(y) = A_x(y - x)$.

Let $p \in S^m(U \times \mathbb{R}^{d+1})$. As $p(x, -iX) = p_\sigma(x, D)$ with $p_\sigma(x, \xi) = p(x, \sigma(x, \xi)) = p(x, (A_x^t)^{-1}\xi)$ the distribution kernel $k_{p(x, -iX)}(x, y)$ of $p(x, -iX)$ is represented by the oscillating integrals

$$(3.26) \quad (2\pi)^{-(d+1)} \int e^{i(x-y) \cdot \xi} p(x, (A_x^t)^{-1}\xi) d\xi = (2\pi)^{-(d+1)} |A_x| \int e^{iA_x(x-y) \cdot \xi} p(x, \xi) d\xi.$$

Since $\psi_x(y) = A_x(y - x)$ we deduce that

$$(3.27) \quad k_{p(x, -iX)}(x, y) = |\psi'_x| \check{p}_{\xi \rightarrow y}(x, -\psi_x(y)).$$

Combining this with Lemma 3.14 then gives:

Proposition 3.15 ([BG, Thms. 15.39, 15.49]). Let $P : C_c^\infty(U) \rightarrow C^\infty(U)$ be a continuous linear operator with distribution kernel $k_P(x, y)$. Then the following are equivalent:

- (i) P is a Ψ_H DO of order m , $m \in \mathbb{C}$.
- (ii) There exist $K \in \mathcal{K}^{\hat{m}}(U \times \mathbb{R}^{d+1})$, $\hat{m} = -(m+d+2)$, and $R \in C^\infty(U \times U)$ such that

$$(3.28) \quad k_P(x, y) = |\psi'_x| K(x, -\psi_x(y)) + R(x, y).$$

Furthermore, if (i) and (ii) hold and $K \sim \sum_{j \geq 0} K_{\hat{m}+j}$, $K_l \in \mathcal{K}_l(U \times \mathbb{R}^{d+1})$, then P has symbol $p \sim \sum_{j \geq 0} p_{m-j}$, $p_l \in S_l(U \times \mathbb{R}^{d+1})$, where p_{m-j} is the restriction to $U \times (\mathbb{R}^{d+1} \setminus 0)$ of $(K_{m+j})_{y \rightarrow \xi}^\wedge$.

In the sequel we will need a version of Proposition 3.15 in Heisenberg coordinates. To this end let ε_x denote the coordinate change to the Heisenberg coordinates at x and set $\phi_x = \varepsilon_x \circ \psi_x^{-1}$. Recall that ϕ_x is a Lie group isomorphism from $G^{(x)}$ to $G_x U$ such that $\phi_x(\lambda \cdot y) = \lambda \cdot \phi_x(y)$ for any $\lambda \in \mathbb{R}$. Moreover, using (2.25) one can check that $|\phi'_x| = 1$ and $\phi_x^{-1}(y) = -\phi_x(-y)$. Therefore, from (3.27) we see that we can put $k_{p(x, -iX)}(x, y)$ into the form

$$(3.29) \quad k_{p(x, -iX)}(x, y) = |\varepsilon'_x| K_P(x, -\varepsilon_x(y)), \quad K_P(x, y) = \check{p}_{\xi \rightarrow y}(x, -\phi_x(-y)) = \check{p}_{\xi \rightarrow y}(x, \phi_x^{-1}(y)).$$

In fact, the coordinate changes ϕ_x give rise to an action on distributions on $U \times \mathbb{R}^{d+1}$ given by

$$(3.30) \quad K(x, y) \longrightarrow \phi_x^* K(x, y), \quad \phi_x^* K(x, y) = K(x, \phi_x^{-1}(y)).$$

Since ϕ_x depends smoothly on x , this action induces a continuous linear isomorphisms of $C^N(U \times \mathbb{R}^{d+1})$, $N \geq 0$, and $C^\infty(U \times \mathbb{R}^{d+1})$ onto themselves. As $\phi_x(y)$ is polynomial in y in such way that $\phi_x(0) = 0$ and $\phi_x(\lambda \cdot y) = \lambda \cdot \phi_x(y)$ for every $\lambda \in \mathbb{R}$, we deduce that the above action also yields a continuous linear isomorphism of $C^\infty(U) \hat{\otimes} \mathcal{S}'_{\text{reg}}(\mathbb{R}^{d+1})$ onto itself and, for every $\lambda > 0$, we have

$$(3.31) \quad (\phi_x^* K)(x, \lambda \cdot y) = \phi_x^* [K(x, \lambda \cdot y)], \quad K \in \mathcal{D}'(U \times \mathbb{R}^{d+1}).$$

Furthermore, as $\phi_x(y)$ is polynomial in y we see that for every $\alpha \in \mathbb{N}^{d+1}$ we can write $\phi_x(y)^\alpha$ in the form $\phi_x(y)^\alpha = \sum_{\langle \beta \rangle = \langle \alpha \rangle} d_{\alpha\beta}(x) y^\beta$ with $d_{\alpha\beta} \in C^\infty(U \times \mathbb{R}^{d+1})$. It then follows that, for every $m \in \mathbb{C}$, the map $K(x, y) \rightarrow \phi_x^* K(x, y)$ induces a linear isomorphisms of $\mathcal{K}_m(U \times \mathbb{R}^{d+1})$ and $\mathcal{K}^m(U \times \mathbb{R}^{d+1})$ onto themselves. Combining this with (3.29) and Proposition 3.15 then gives:

Proposition 3.16. *Let $P : C_c^\infty(U) \rightarrow C^\infty(U)$ be a continuous linear operator with distribution kernel $k_P(x, y)$. Then the following are equivalent:*

(i) *P is a Ψ_H DO of order m , $m \in \mathbb{C}$.*

(ii) *There exist $K_P \in \mathcal{K}^{\hat{m}}(U \times \mathbb{R}^{d+1})$, $\hat{m} = -(m + d + 2)$, and $R \in C^\infty(U \times U)$ such that*

$$(3.32) \quad k_P(x, y) = |\varepsilon'_x| K_P(x, -\varepsilon_x(y)) + R(x, y).$$

Furthermore, if (i) and (ii) hold and $K_P \sim \sum_{j \geq 0} K_{P, \hat{m}+j}$, $K_l \in \mathcal{K}_l(U \times \mathbb{R}^{d+1})$, then P has symbol $p \sim \sum_{j \geq 0} p_{m-j}$, $p_l \in S_l(U \times \mathbb{R}^{d+1})$, where p_{m-j} is the restriction to $U \times (\mathbb{R}^{d+1} \setminus 0)$ of $[K_{P, \hat{m}+j}(x, \phi_x^{-1}(y))]_{y \rightarrow \xi}^\wedge$.

Remark 3.17. Let $a \in U$. Then (3.32) shows that the distribution kernel of $\tilde{P} = (\varepsilon_a)_* P$ at $x = 0$ is

$$(3.33) \quad k_{\tilde{P}}(0, y) = |\varepsilon'_a|^{-1} k_P(\varepsilon_a^{-1}(0), \varepsilon_a^{-1}(y)) = K_P(a, -y).$$

Moreover, as we are in Heisenberg coordinates already, we have $\psi_0 = \varepsilon_0 = \phi_0 = \text{id}$. Thus, in the form (3.32) for $\tilde{P}$ we have $K_{\tilde{P}}(0, y) = K_P(a, y)$. Therefore, if we let $p_m(x, \xi)$ denote the principal symbol of P and let $K_{P, \hat{m}} \in \mathcal{K}_{\hat{m}}(U \times \mathbb{R}^{d+1})$ denote the leading kernel of K_P , then by Proposition 3.16 we have

$$(3.34) \quad p_m(0, \xi) = [K_{P, \hat{m}}]_{y \rightarrow \xi}^\wedge(a, \xi).$$

This shows that $[K_{P, \hat{m}}]_{y \rightarrow \xi}^\wedge(a, \xi)$ is the principal symbol of P at $x = 0$ in Heisenberg coordinates centered at a .

3.5. Ψ_H DO's on a general Heisenberg manifold. Let (M^{d+1}, H) be a Heisenberg manifold. As alluded to before the Ψ_H DO's on an subset of $\mathbb{R}^{d+1}$ are Ψ DO's of type $(\frac{1}{2}, \frac{1}{2})$. However, the latter don't make sense on a general manifold, for their class is not preserved by an arbitrary change of chart. Nevertheless, when dealing with Ψ_H DO's this issue is resolved if we restrict ourselves to changes of Heisenberg charts. Indeed, we have:

Proposition 3.18. *Let U (resp. $\tilde{U}$) be an open subset of $\mathbb{R}^{d+1}$ together with a hyperplane bundle $H \subset TU$ (resp. $\tilde{H} \subset T\tilde{U}$) and a H -frame of TU (resp. a $\tilde{H}$ -frame of $T\tilde{U}$). Let $\phi : (U, H) \rightarrow (\tilde{U}, \tilde{H})$ be a Heisenberg diffeomorphism and let $\tilde{P} \in \Psi_{\tilde{H}}^m(\tilde{U})$.*

1) *The operator $P = \phi^* \tilde{P}$ is a Ψ_H DO of order m on U .*

2) *If the distribution kernel of $\tilde{P}$ is of the form (3.32) with $K_{\tilde{P}}(\tilde{x}, \tilde{y}) \in \mathcal{K}^{\hat{m}}(\tilde{U} \times \mathbb{R}^{d+1})$ then the distribution kernel of P can be written in the form (3.32) with $K_P(x, y) \in \mathcal{K}^{\hat{m}}(U \times \mathbb{R}^{d+1})$ such that*

$$(3.35) \quad K_P(x, y) \sim \sum_{\langle \beta \rangle \geq \frac{3}{2} \langle \alpha \rangle} \frac{1}{\alpha! \beta!} a_{\alpha\beta}(x) y^\beta (\partial_y^\alpha K_{\tilde{P}})(\phi(x), \phi'_H(x)y),$$

where we have let $a_{\alpha\beta}(x) = \partial_y^\beta [|\partial_y(\tilde{\varepsilon}_{\phi(x)} \circ \phi \circ \varepsilon_x^{-1})(y)|(\tilde{\varepsilon}_{\phi(x)} \circ \phi \circ \varepsilon_x^{-1}(y) - \phi'_H(x)y)^\alpha]_{|y=0}$ and $\tilde{\varepsilon}_{\tilde{x}}$ denote the change to the Heisenberg coordinates at $\tilde{x} \in \tilde{U}$. In particular, we have

$$(3.36) \quad K_P(x, y) = |\phi'_H(x)| K_{\tilde{P}}(\phi(x), \phi'_H(x)y) \mod \mathcal{K}^{\hat{m}+1}(U \times \mathbb{R}^{d+1}).$$

Remark 3.19. The version of the above statement in [BG] does not contain the asymptotics (3.36), which will be crucial for giving a global definition of the principal symbol of a Ψ_H DO in the next section. For this reason give a detailed proof of the above version in Section 8. This proof will also be useful in [Po3] and [Po4] for generalizing Proposition 3.18 to holomorphic families of Ψ_H DO's and to Ψ_H DO's with parameter.

As a consequence of Proposition 3.18 we can define Ψ_H DO's on M acting on the sections of a vector bundle $\mathcal{E}$ over M .

Definition 3.20. $\Psi_H^m(M, \mathcal{E})$, $m \in \mathbb{C}$, consists of continuous operators $P : C_c^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ such that:

- (i) *The distribution kernel of P is smooth off the diagonal;*
- (ii) *For any trivialization $\tau : \mathcal{E}|_U \rightarrow U \times \mathbb{C}^r$ over a local Heisenberg chart $\kappa : U \rightarrow V \subset \mathbb{R}^{d+1}$ the operator $\kappa_* \tau_*(P|_U)$ belongs to $\Psi_H^m(V, \mathbb{C}^r) := \Psi_H^m(V) \otimes \text{End } \mathbb{C}^r$.*

All the previous properties of Ψ DO's on an open subset of $\mathbb{R}^{d+1}$ hold *mutatis standis* for Ψ_H DO's on M acting on sections of $\mathcal{E}$.

3.6. Transposes and adjoints of Ψ_H DO's. Let us now look at the transpose and adjoints of Ψ_H DO's. First, given a Heisenberg chart $U \subset \mathbb{R}^{d+1}$ we have:

Proposition 3.21. *Let $P \in \Psi_H^m(U)$. Then:*

- 1) *The transpose operator P^t is a Ψ_H DO of order m on U .*
- 2) *If we write the distribution kernel of P in the form (3.32) with $K_P \in \mathcal{K}^{\hat{m}}(U \times \mathbb{R}^{d+1})$ then P^t can be written in the form (3.32) with $K_{P^t} \in \mathcal{K}^{\hat{m}}(U \times \mathbb{R}^{d+1})$ such that*

$$(3.37) \quad K_{P^t}(x, y) \sim \sum_{\frac{3}{2} \langle \alpha \rangle \leq \langle \beta \rangle} \sum_{|\gamma| \leq |\delta| \leq 2|\gamma|} a_{\alpha\beta\gamma\delta}(x) y^{\beta+\delta} (\partial_x^\gamma \partial_y^\alpha K_P)(x, -y),$$

where $a_{\alpha\beta\gamma\delta}(x) = \frac{|\varepsilon_x^{-1}|}{\alpha! \beta! \gamma! \delta!} [\partial_y^\beta (|\varepsilon'_{\varepsilon_x^{-1}(-y)}| (y - \varepsilon_{\varepsilon_x^{-1}(y)}(x))^\alpha) \partial_y^\delta (\varepsilon_x^{-1}(-y) - x)^\gamma](x, 0)$. In particular,

$$(3.38) \quad K_{P^t}(x, y) = K_P(x, -y) \mod \mathcal{K}^{\hat{m}+1}(U \times \mathbb{R}^{d+1}).$$

Remark 3.22. The asymptotic expansion (3.37) is not stated in [BG], but we need it in order to determine the global principal symbol of the transpose of a Ψ_H DO (see next section). A detailed proof of Proposition 3.21 can be found in Section 9.

Using this result, or its version in [BG], we obtain:

Proposition 3.23 ([BG, Thm. 17.4]). *Let $P : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a Ψ_H DO of order m . Then:*

- 1) *The transpose operator $P^t : \mathcal{E}'(M, \mathcal{E}^*) \rightarrow \mathcal{D}'(M, \mathcal{E}^*)$ is a Ψ_H DO of order m ;*
- 2) *If M is endowed with a smooth positive density and $\mathcal{E}$ with a Hermitian metric then the adjoint operator $P^* : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ is a Ψ_H DO of order $\overline{m}$.*

4. PRINCIPAL SYMBOL AND MODEL OPERATORS.

In this section we define the principal symbols and model operators of Ψ_H DO's and check their main properties.

4.1. Principal symbol and model operators. Let $\mathfrak{g}^*M$ be the dual bundle of $\mathfrak{g}M$ with canonical projection $\pi : \mathfrak{g}^*M \rightarrow M$.

Definition 4.1. $S_m(\mathfrak{g}^*M, \mathcal{E})$, $m \in \mathbb{C}$, is the space of sections $p \in C^\infty(\mathfrak{g}^*M \setminus 0, \text{End } \pi^*\mathcal{E})$ which are homogeneous of degree m in the sense that, for any $\lambda > 0$, we have

$$(4.1) \quad p(x, \lambda\xi) = \lambda^m p(x, \xi) \quad \forall (x, \xi) \in \mathfrak{g}^*M \setminus 0,$$

where $\xi \rightarrow \lambda\xi$ denotes the dilation (3.2).

Let $P \in \Psi_H^m(M)$ and for $j = 1, 2$ let κ_j be a Heisenberg chart with domain $V_j \subset M$ and let $\phi : U_1 \rightarrow U_2$ be the corresponding transition map, where we have let $U_j = \kappa_j(V_1 \cap V_2) \subset \mathbb{R}^{d+1}$.

Let us first assume that $\mathcal{E}$ is the trivial line bundle, so that P is a scalar operator. For $j = 1, 2$ we let $P_j := \kappa_{j*}(P|_{V_1 \cap V_2})$, so that $P_1 = \phi^*P_2$. Since P_j belongs to $\Psi_H^m(U_j)$ its distribution kernel is of the form (3.32) with $K_{P_j} \in \mathcal{K}^{\hat{m}}(U_j \times \mathbb{R}^{d+1})$. Moreover, by Proposition 3.18 we have

$$(4.2) \quad K_{P_1}(x, y) = |\phi'_H(x)| K_{P_2}(\phi(x), \phi'_H(x)y) \mod \mathcal{K}^{\hat{m}+1}(U_1 \times \mathbb{R}^{d+1}).$$

Therefore, if we let $K_{P_j, \hat{m}} \in \mathcal{K}^{\hat{m}}(U_j \times \mathbb{R}^{d+1})$ be the leading kernel of K_{P_j} then we get

$$(4.3) \quad K_{P_1, \hat{m}}(x, y) = |\phi'_H(x)| K_{P_2, \hat{m}}(\phi(x), \phi'_H(x)y).$$

Next, for $j = 1, 2$ we define

$$(4.4) \quad p_{j, m}(x, \xi) = [K_{P_j, \hat{m}}]_{y \rightarrow \xi}^\wedge(x, \xi), \quad (x, \xi) \in U_j \times \mathbb{R}^{d+1} \setminus 0.$$

By Remark 3.17 for any $a \in U_j$ the symbol $p_j(a, \cdot)$ yields in Heisenberg coordinates centered at a the principal symbol of P_j at $x = 0$. Moreover, since $\phi'_H(a)$ is a linear map, from (4.3) we get

$$(4.5) \quad p_{1, m}(x, \xi) = p_{2, m}(\phi(x), [\phi'_H(x)]^{-1t}\xi).$$

This shows that $p_m := \kappa_1^* p_{1, m}$ is an element of $S_m(\mathfrak{g}^*(V_1 \cap V_1))$ which is independent of the choice of the chart κ_1 . Since $S_m(\mathfrak{g}^*M)$ is a sheaf this gives rise to a uniquely defined symbol $p_m \in S_m(\mathfrak{g}^*M)$.

When $\mathcal{E}$ is a general vector bundle, the above construction can be carried out similarly, so that we obtain:

Theorem 4.2. *For any $P \in \Psi_H^m(M, \mathcal{E})$ there is a unique symbol $\sigma_m(P) \in S_m(\mathfrak{g}^*M, \mathcal{E})$ such that if in a local trivializing Heisenberg chart $U \subset \mathbb{R}^{d+1}$ we let $K_{P, \hat{m}}(x, y) \in \mathcal{K}^{\hat{m}}(U \times \mathbb{R}^{d+1})$ be the leading kernel for the kernel $K_P(x, y)$ in the form (3.32) for P , then we have*

$$(4.6) \quad \sigma_m(P)(x, \xi) = [K_{P, \hat{m}}]_{y \rightarrow \xi}^\wedge(x, \xi), \quad (x, \xi) \in U \times \mathbb{R}^{d+1} \setminus 0.$$

Equivalently, for any $x_0 \in M$ the symbol $\sigma_m(P)(x_0, \cdot)$ agrees in trivializing Heisenberg coordinates centered at x_0 with the principal symbol of P at $x = 0$.

Definition 4.3. *For $P \in \Psi_H^m(M, \mathcal{E})$ the symbol $\sigma_m(P) \in S_m(\mathfrak{g}^*M, \mathcal{E})$ provided by Theorem 4.2 is called the principal symbol of P .*

Remark 4.4. Since we have two notions of principal symbol we shall distinguish between them by saying that $\sigma_m(P)$ is the global principal symbol of P and that in a local trivializing chart the principal symbol p_m of P in the sense of (3.13) is the local principal symbol of P in this chart.

In a local Heisenberg chart $U \subset \mathbb{R}^{d+1}$ the global symbol $\sigma_m(P)$ and the local principal symbol p_m of $P \in \Psi_H^m(U)$ can be easily related to each other. Indeed, by Proposition 3.16 we have

$$(4.7) \quad p_m(x, \xi) = [K_{P, \hat{m}}(x, \phi_x^{-1}(y))]_{y \rightarrow \xi}^\wedge(x, \xi),$$

where $K_{P, \hat{m}}$ denotes the leading kernel for the kernel K_P in the form (3.32) for P . By combining this with the definition (4.4) of $\sigma_m(P)$ we thus get

$$(4.8) \quad p_m(x, \xi) = (\hat{\phi}_x^* \sigma_m(P))(x, \xi),$$

$$(4.9) \quad (\hat{\phi}_x^* \sigma_m(P)) = [[\sigma_m(P)]_{\xi \rightarrow y}^\vee(x, \phi_x^{-1}(y))]_{y \rightarrow \xi}^\wedge = [\phi_x^* [\sigma_m(P)]_{\xi \rightarrow y}^\vee]_{y \rightarrow \xi}^\wedge,$$

where ϕ_x^* is the isomorphism map (3.30). In particular, since the latter is a linear isomorphism of $\mathcal{K}_m(U \times \mathbb{R}^{d+1})$ onto itself, we see that the map $p \rightarrow \hat{\phi}_x^* p$ is a linear isomorphism of $S_m(U \times \mathbb{R}^{d+1})$ onto itself.

Example 4.5. Let $X_0, \dots, X_d$ be a local H -frame of TM near a point $a \in M$. In any Heisenberg chart associated with this frame the Heisenberg symbol of X_j is $\frac{1}{i}\xi_j$. In particular, this is true in Heisenberg coordinates centered at a . Thus the (global) principal symbol of X_j is equal to $\frac{1}{i}\xi_j$ in the local trivialization of $\mathfrak{g}^*M \setminus 0$ defined by the frame $X_0, \dots, X_d$. More generally, for any differential $P = \sum_{\langle \alpha \rangle \leq m} a_\alpha(x) X^\alpha$ on M we have

$$(4.10) \quad \sigma_m(P)(x, \xi) = \sum_{\langle \alpha \rangle \leq m} a_\alpha(x) i^{-|\alpha|} \xi^\alpha.$$

Thus, for differential operators the global and local principal symbols agree in suitable coordinates. Alternatively, this result follows from the fact that the isomorphism (3.30) induces the identity map on distributions $K(x, y)$ supported in $U \times \{y = 0\}$.

Proposition 4.6. *For every $m \in \mathbb{C}$ the principal symbol map $\sigma_m : \Psi_H^m(M, \mathcal{E}) \rightarrow S_m(\mathfrak{g}^*M, \mathcal{E})$ gives rise to a linear isomorphism $\Psi_H^m(M, \mathcal{E})/\Psi_H^{m-1}(M, \mathcal{E}) \xrightarrow{\sim} S_m(\mathfrak{g}^*M, \mathcal{E})$.*

Proof. By construction the principal symbol of $P \in \Psi_H^m(M, \mathcal{E})$ depends only on his principal part in local coordinates and vanishes everywhere if, and only if, the order of P is $\leq m-1$. Therefore, the kernel of the principal symbol map σ_m is $\Psi_H^{m-1}(M, \mathcal{E})$, so σ_m induces an injective linear map $\Psi_H^m(M, \mathcal{E})/\Psi_H^{m-1}(M, \mathcal{E}) \rightarrow S_m(\mathfrak{g}^*M, \mathcal{E})$.

To complete the proof it is enough to show that σ_m is surjective. To this end consider a symbol $p_m(x, \xi) \in S_m(\mathfrak{g}^*M, \mathcal{E})$ and let $(\varphi_i)_{i \in I}$ be a partition of the unity subordinated to an open covering $(U_i)_{i \in I}$ of M by domains of Heisenberg charts $\kappa_i : U_i \rightarrow V_i$ over which there are trivializations $\tau_i : \mathcal{E}|_{U_i} \rightarrow U_i \times \mathbb{C}^r$. For each index i let $\psi_i \in C^\infty(U_i)$ be such that $\psi_i = 1$ near $\text{supp } \varphi_i$ and set

$$(4.11) \quad p_m^{(i)}(x, \xi) = (1 - \chi(\xi))(\hat{\phi}_{i,x}^* \kappa_{i*} \tau_{i*} p_m|_{\mathfrak{g}^*U_i \setminus 0})(x, \xi) \in S_m(V_i) \otimes \text{End } \mathbb{C}^r,$$

where $\chi \in C^\infty(\mathbb{R}^{d+1})$ is such that $\chi = 1$ near the origin and $\hat{\phi}_{i,x}^*$ denotes the isomorphism (4.9) with respect to the chart V_i . Then we define a Ψ_H DO of order m by letting

$$(4.12) \quad P = \sum \varphi_i [\tau_i^* \kappa_i^* p_m^{(i)}(x, -iX)] \psi_i.$$

For for every index i the local principal symbol of $\varphi_i [\tau_i^* \kappa_i^* p_m^{(i)}(x, -iX)] \psi_i$ in the chart V_i is $\varphi_i \circ \kappa_i^{-1}(\hat{\phi}_{i,x}^* \kappa_{i*} \tau_{i*} p_m|_{\mathfrak{g}^*U_i \setminus 0})$, so by (4.8) its global principal is $\varphi_i \circ \kappa_i^{-1}(\kappa_{i*} \tau_{i*} p_m|_{\mathfrak{g}^*U_i \setminus 0})$, which pulls back to $\varphi_i p_m$ on U_i . It follows that the global principal symbol of P is $\sigma_m(P) = \sum_i \varphi_i p_m = p_m$. This proves the surjectivity of the map σ_m , so the proof is now complete. $\square$

Next, granted the above definition of the principal symbol, we can define the model operator at a point as follows.

Definition 4.7. Let $P \in \Psi_H^m(M, \mathcal{E})$ have (global) principal symbol $\sigma_m(P)$. Then the model operator of P at $a \in M$ is the left-invariant Ψ_H DO-operator $P^a : \mathcal{S}_0(G_a M, \mathcal{E}_a) \rightarrow \mathcal{S}_0(G_a M, \mathcal{E}_a)$ with symbol $\sigma_m(P)_{\xi \rightarrow y}^\vee(a, \cdot)$, i.e.,

$$(4.13) \quad P^a f(x) = \langle \sigma_m(P)_{\xi \rightarrow y}^\vee(a, y), f(x \cdot y^{-1}) \rangle, \quad f \in \mathcal{S}_0(G_a M, \mathcal{E}_a).$$

Consider a local trivializing chart $U \subset \mathbb{R}^{d+1}$ near a and let us relate the model operator P^a on $G_a M$ to the operator $P^{(a)} = \tilde{p}_m^a(-iX^{(a)})$ on $G^{(a)}$ defined using the local principal symbol $\tilde{p}_m(x, \xi)$ of P in this chart. Using (3.6) and (4.8) for $f \in \mathcal{S}_0(\mathbb{R}^{d+1})$ we get

$$(4.14) \quad P^{(a)} f(y) = \langle (p_m^a)^\vee(z), f(y \cdot z^{-1}) \rangle = \langle (\sigma_m(P)_{\xi \rightarrow y}^\vee(x, \phi_a^{-1}(y))), f(y \cdot z^{-1}) \rangle.$$

Since $|\phi'_a| = 1$ and ϕ_a is a Lie group isomorphism from $G^{(a)}$ onto $G_a M$ we obtain

$$(4.15) \quad P^{(a)} f(y) = \langle (\sigma_m(P)_{\xi \rightarrow y}^\vee(x, y), f \circ \phi_a^{-1}(y \cdot \phi_a(x)^{-1}) \rangle = (\phi_a^* P^a) f(y).$$

In particular, we have

$$(4.16) \quad P^a = (\phi_a)_* p_m^a(-iX^{(a)}).$$

4.2. Composition of principal symbols and model operators. Let us now look at the composition of principal symbols. To this end for $a \in M$ we let $*^a : S_{m_1}(\mathbb{R}^{d+1}) \times S_{m_2}(\mathbb{R}^{d+1}) \rightarrow S_{m_1+m_2}(\mathbb{R}^{d+1})$ be the convolution product for symbols defined by the product law of $G_a M$ under the identification $G_a M \simeq \mathbb{R}^{d+1}$ provided by a H -frame $X_0, \dots, X_d$ of TM near a , that is,

$$(4.17) \quad (p_{m_1} *^a p_{m_2})(-iX^a) = p_{m_1}(-iX^a) \circ p_{m_2}(-iX^a), \quad p_{m_j} \in S_{m_j}(\mathbb{R}^{d+1}).$$

Let $U \subset \mathbb{R}^{d+1}$ be a local trivializing Heisenberg chart near a and for $j = 1, 2$ let $P_j \in \Psi_H^{m_j}(U)$ have (global) principal symbol $\sigma_{m_j}(P_j)$. Recall that under the trivialization of GU provided by the H -frame $X_0, \dots, X_d$ we have $P_j^a = \sigma(P_j)(x, -iX^a)$. Thus,

$$(4.18) \quad [\sigma_{m_j}(P_j)(x, \cdot) *^a \sigma_{m_j}(P_j)(x, \cdot)](-iX^a) = P_1^a P_2^a.$$

On the other hand, using (4.8) and (4.16) we see that $\hat{\phi}_a^*[p_{m_1} *^a p_{m_2}](-iX^a)$ is equal to

$$(4.19) \quad \begin{aligned} \hat{\phi}_a^*[p_{m_1}(-iX^a) \circ p_{m_2}(-iX^a)] &= \hat{\phi}_a^*[p_{m_1}(-iX^a)] \circ \hat{\phi}_a^*[p_{m_2}(-iX^a)] \\ &= (\hat{\phi}_a^* p_{m_1})(-iX^{(a)}) \circ (\hat{\phi}_a^* p_{m_2})(-iX^{(a)}) = [(\hat{\phi}_a^* p_{m_1}) *^{(a)} (\hat{\phi}_a^* p_{m_2})](-iX^{(a)}). \end{aligned}$$

Hence we have

$$(4.20) \quad p_{m_1} *^a p_{m_2} = (\hat{\phi}_a)_* [(\hat{\phi}_a^* p_{m_1}) *^{(a)} (\hat{\phi}_a^* p_{m_2})] \quad \forall p_{m_j} \in S_{m_j}(\mathbb{R}^{d+1}),$$

where $(\hat{\phi}_a)_*$ denotes the inverse of $\hat{\phi}_a^*$. Since $\hat{\phi}_a^*$, its inverse and $*^{(a)}$ depend smoothly on a , we deduce that so does $*^a$. Therefore, we obtain:

Proposition 4.8. The group laws on the fibers of GM give rise to a convolution product,

$$(4.21) \quad * : S_{m_1}(\mathfrak{g}^* M, \mathcal{E}) \times S_{m_2}(\mathfrak{g}^* M, \mathcal{E}) \longrightarrow S_{m_1+m_2}(\mathfrak{g}^* M, \mathcal{E}),$$

$$(4.22) \quad p_{m_1} * p_{m_2}(x, \xi) = [p_{m_1}(x, \cdot) *^x p_{m_2}(x, \cdot)](\xi), \quad p_{m_j} \in S_{m_j}(\mathfrak{g}^* M, \mathcal{E}),$$

where $*^x$ denote the convolution product for symbols on $G_x M$.

Notice that (4.20) shows that, under the relation (4.8) between local and global principal symbols, the convolution product (4.17) for global principal symbols corresponds to the convolution product (3.19) for local principal symbols. Since by Proposition 3.9 the latter yields the *local* principal symbol of the product of two Ψ_H DO's in a local chart, we deduce that the convolution product (4.17) yields the *global* principal symbol of the product two Ψ_H DO's.

Moreover, by (4.18) the global convolution product (4.17) corresponds to the product of model operators, so the model operator of a product of two $\Psi_H\text{DO}$'s is equal to the product of the model operators. We have thus proved:

Proposition 4.9. *For $j = 1, 2$ let $P_j \in \Psi_H^{m_j}(M, \mathcal{E})$ and suppose that P_1 or P_2 is properly supported.*

- 1) *We have $\sigma_{m_1+m_2}(P_1 P_2) = \sigma_{m_1}(P) * \sigma_{m_2}(P)$.*
- 2) *At every $a \in M$ the model operator of $P_1 P_2$ is $(P_1 P_2)^a = P_1^a P_2^a$.*

4.3. Principal symbol of transposes and adjoints. In this subsection we shall determine the principal symbols and the model operators of transposes and adjoints of $\Psi_H\text{DO}$'s.

Recall that if $P \in \Psi_H^m(M, \mathcal{E})$ then by Proposition 3.23 its transpose $P^t : C_c^\infty(M, \mathcal{E}^*) \rightarrow C^\infty(M, \mathcal{E})$ is a $\Psi_H\text{DO}$ of order m and its adjoint $P^* : C_c^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ is a $\Psi_H\text{DO}$ of order $\bar{m}$ (assuming M endowed with a positive density and $\mathcal{E}$ with a Hermitian metric in order to define the adjoint).

Proposition 4.10. *Let $P \in \Psi_H^m(M, \mathcal{E})$ have principal symbol $\sigma_m(P)$. Then:*

- 1) *The principal symbol of P^t is $\sigma_m(P^t)(x, \xi) = \sigma_m(x, -\xi)^t$ (this is an element of $S_m(\mathfrak{g}^*M, \mathcal{E}^*)$);*
- 2) *If P^a is the model operator of P at a then the model operator of P^t at a is the transpose operator $(P^a)^t : S_0(G_x M, \mathcal{E}_x^*) \rightarrow S_0(G_x M, \mathcal{E}_x^*)$.*

Proof. Let us first assume that $\mathcal{E}$ is the trivial line bundle and that P is a scalar operator. In a local Heisenberg chart $U \subset \mathbb{R}^{d+1}$ we can write the distribution kernels of P and P^t in the form (3.32) with K_P and K_{P^t} in $\mathcal{K}^{\hat{m}}(U \times \mathbb{R}^{d+1})$. Let $K_{P, \hat{m}}$ and $K_{P^t, \hat{m}}^t$ denote the principal parts of K_P and K_{P^t} respectively. Then the principal symbols of P and P^t are $\sigma_m(P)(x, \xi) = (K_{P, \hat{m}})_{y \rightarrow \xi}^\wedge(x, \xi)$ and $\sigma_m(P^t)(x, \xi) = (K_{P^t, \hat{m}})_{y \rightarrow \xi}^\wedge(x, \xi)$ respectively. Since (3.38) implies that $K_{P^t, \hat{m}}(x, y) = K_{P, \hat{m}}(x, -y)$ and the Fourier transform commutes with the multiplication by -1 we get

$$(4.23) \quad \sigma_m(P^t)(x, \xi) = \sigma_m(P)(x, -\xi).$$

Next, for $a \in U$ let $p \in S_m(G_a)$ and let P be the left-invariant $\Psi_H\text{DO}$ with symbol p . Then the transpose P^t is such that, for f and g in $S_0(G_a U)$, we have

$$(4.24) \quad \langle P^t f, g \rangle = \langle f, P v \rangle = \langle 1, f(x)(P g)(x) \rangle = \langle 1, f(x) \langle \check{p}(y), g(x.y^{-1}) \rangle \rangle \\ = \langle 1 \otimes \check{p}(x, y), f(x) g(x.y^{-1}) \rangle.$$

Therefore, using the change of variable $(x, y) \rightarrow (x.y^{-1}, y^{-1})$ and the fact $y^{-1} = -y$ we get

$$(4.25) \quad \langle P^t f, g \rangle = \langle 1 \otimes \check{p}(x, -y), f(x) g(x.y^{-1}) \rangle = \langle 1, f(x) \langle \check{p}(-y), g(x.y^{-1}) \rangle \rangle.$$

Since $\check{p}(-y) = \check{p}^t(y)$ with $p^t(\xi) = p(-\xi)$, we obtain

$$(4.26) \quad \langle P^t f, g \rangle = \langle 1, f(x) \langle \check{p}^t(y), g(x.y^{-1}) \rangle \rangle = \langle (p^t * f)(x), g(x) \rangle.$$

Thus P^t is the left-convolution operator with symbol $p^t(\xi) = p(-\xi)$.

Now, since the model operator $(P^t)^a$ is the left-invariant $\Psi_H\text{DO}$ with symbol $\sigma_m(P^t)(a, \xi) = \sigma_m(P)(a, -\xi)$, it follows that $(P^t)^a$ agrees with the transpose of P^a .

In the general case, when $\mathcal{E}$ is not the trivial bundle, we can similarly show that P^t is a $\Psi_H\text{DO}$ of order m with principal symbol $\sigma_m(P^t)(x, \xi) = \sigma_m(P)(x, -\xi)^t$ and such that at every point $a \in M$ its model operator at a is the transpose $(P^a)^t$ of P^a . $\square$

Assume now that M is endowed with a positive density and $\mathcal{E}$ with a Hermitian metric respectively and let $L^2(M, \mathcal{E})$ be the associated L^2 -Hilbert space.

Proposition 4.11. *Let $P \in \Psi_H^m(M, \mathcal{E})$ have principal symbol $\sigma_m(P)$. Then:*

- 1) *The principal symbol of P^* is $\sigma_{\bar{m}}(P^*)(x, \xi) = \sigma_m(P)(x, \xi)^*$.*
- 2) *If P^x denotes the model operator of P at $x \in M$ then the model operator of P^* at x is the adjoint $(P^x)^*$ of P^x .*

Proof. Let us first assume that $\mathcal{E}$ is the trivial line bundle, so that P is a scalar operator. Moreover, since the above statements are local ones, it is enough to prove them in a local Heisenberg chart $U \subset \mathbb{R}^{d+1}$ and we may assume that P is a Ψ_H DO on U .

Let $\bar{P} : C_c^\infty(U) \rightarrow C^\infty(U)$ be the conjugate operator of P , so that $\bar{P}u = \overline{P(\bar{f})}$ for any $f \in C_c^\infty(U)$. By Proposition 3.16 the distribution kernel of P of the form (3.32) with $K_P(x, y)$ in $\mathcal{K}_{\hat{m}}(U \times \mathbb{R}^{d+1})$, so the kernel of $\bar{P}$ takes the form

$$(4.27) \quad k_{\bar{P}}(x, y) = \overline{k_P(x, y)} = |\varepsilon'_x| \overline{K_P(x, y)} \mod C^\infty(U \times U).$$

Since the conjugation of distribution $K(x, y) \rightarrow \overline{K(x, y)}$ induces an anti-linear isomorphism from $\mathcal{K}^{\hat{m}}(U \times \mathbb{R}^{d+1})$ onto $\mathcal{K}^{\hat{m}}(U \times \mathbb{R}^{d+1})$, it follows from Proposition 3.16 that $\bar{P}$ is a Ψ_H DO of order $\hat{m}$ and its kernel can be put into the form (3.32) with $K_{\bar{P}}(x, y) = \overline{K_P(x, y)}$. In particular, if we let $K_{P, \hat{m}} \in \mathcal{K}_{\hat{m}}(U \times \mathbb{R}^{d+1})$ denote the leading kernel of K_P then the leading kernel of $K_{\bar{P}}$ is $\overline{K_{P, \hat{m}}}$. Thus $\bar{P}$ has principal symbol,

$$(4.28) \quad \sigma_{\bar{m}}(\bar{P})(x, \xi) = \overline{[K_{P, \hat{m}}]_{\xi \rightarrow y}^\wedge(x, \xi)} = \overline{[(K_{P, \hat{m}})_{\xi \rightarrow y}^\wedge(x, -\xi)]} = \overline{\sigma_m(x, -\xi)}.$$

Moreover, since $\sigma_{\bar{m}}(\bar{P})_{\xi \rightarrow y}^\vee(x, y) = \overline{\sigma_m(P)_{\xi \rightarrow y}^\vee(x, y)}$ the model operator at $a \in U$ of $\bar{P}$ is such that, for any $f \in \mathcal{S}_0(G_a U)$, we have

$$(4.29) \quad (\bar{P})^a f(x) = \langle \sigma_m(P)_{\xi \rightarrow y}^\vee(x, y), f(x \cdot y^{-1}) \rangle = \overline{\langle \sigma_m(P)_{\xi \rightarrow y}^\vee(x, y), \overline{f(x \cdot y^{-1})} \rangle} = \overline{P^a f(x)},$$

so $(\bar{P})^a$ agrees with $\overline{P^a}$.

Combining this with Proposition 3.21 and Proposition 4.10 we thus see that $\bar{P}^t$ is a Ψ_H DO of order $\bar{m}$ such that:

- If we put the kernel of $\bar{P}^t$ into the form (3.32) with respect to $K_{\bar{P}^t}(x, y) \in \mathcal{K}_{\hat{m}}(U \times \mathbb{R}^{d+1})$, then the leading kernel of $K_{\bar{P}^t}$ is $\overline{K_{P, \hat{m}}(x, -y)}$;
- The global principal symbol of $\bar{P}^t$ is $\sigma_{\bar{m}}(\bar{P}^t) = \overline{\sigma_m(P^t)(x, \xi)} = \overline{\sigma_m(P)(x, \xi)}$;
- The model operator at $a \in U$ of $\bar{P}^t$ is $(\bar{P}^t)^a = \overline{P^a} = (P^a)^*$.

Now, let $d\rho(x) = \rho(x)dx$ be the smooth positive density on U coming from that of M . The formal adjoint $P^* : C_c^\infty(U) \rightarrow C^\infty(U)$ of P with respect to $d\rho$ is such that

$$(4.30) \quad \int_f \overline{P f(x)} g(x) \rho(x) dx = \int_U \overline{f(x)} P^* g(x) \rho(x) dx, \quad f, g \in C_c^\infty(U).$$

Thus $P^* = \rho^{-1} \bar{P}^t \rho$, which shows that P^* is a Ψ_H DO of order $\bar{m}$. Moreover, as in the proof of Proposition 3.21 in Section 9, we can prove that the kernel of P^* can be put into the form (3.32) with $K_{P^*}(x, y) \in \mathcal{K}^{\bar{m}}(U \times \mathbb{R}^{d+1})$ such that

$$(4.31) \quad K_{P^*}(x, y) = \rho(x)^{-1} K_{\bar{P}^t}(x, y) \rho(\varepsilon_x^{-1}(y)) \sim \sum_{\alpha} \frac{1}{\alpha!} \rho(x)^{-1} \partial_y (\rho(\varepsilon_x^{-1}(y)))|_{y=0} K_{\bar{P}^t}(x, y).$$

In particular, the kernels $K_{P^*}(x, y)$ and $K_{\bar{P}^t}(x, y)$ agree modulo $\mathcal{K}^{\hat{m}+1}(U \times \mathbb{R}^{d+1})$, hence have same leading kernel. It then follows that P^* and $\bar{P}^t$ have same principal symbol and same model operator at a point $a \in U$, that is, we have $\sigma_{\bar{m}}(P^*)(a, \xi) = \overline{\sigma_m(P)(a, \xi)}$ and $(P^*)^a = (P^a)^*$.

Finally, assume that $\mathcal{E}$ is a general bundle, so that the restriction of P to U is given by a matrix $P = (P_{ij})$ of Ψ_H DO's of order r . Let $h(x) \in C^\infty(U, GL_r(\mathbb{C}))$, $h(x)^* = h(x)$, be the Hermitian metric on $U \times \mathbb{C}^r$ coming from that of $\mathcal{E}$. Then the adjoint of P with respect to this Hermitian metric is $P^* = \rho^{-1} h^{-1} \bar{P}^t h \rho$. Therefore, in the same way as in the scalar case we can prove that P^* has principal symbol $h(x)^{-1} \overline{\sigma_m(P)(x, \xi)}^t h(x) = \sigma_m(P)(x, \xi)^*$ and its model operator at any point $a \in U$ is $h(a)^{-1} \overline{P^a}^t h(a) = (P^a)^*$. $\square$

5. HYPOELLIPTICITY, PARAMETRICES AND THE ROCKLAND CONDITION

In this section we define a Rockland condition for Ψ_H DO's and relate it to the invertibility of the principal symbol to get hypoellipticity criterions.

First, by [BG, Sect. 18] in a local Heisenberg chart the invertibility of the local principal symbol of a Ψ_H DO is equivalent to the existence of a Ψ_H DO-parametrix. Using the global principal symbol we can give a global reformulation of this result as follows.

Proposition 5.1. *Let $P : C_c^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a Ψ_H DO of order m such that $k := \Re m > 0$. The following are equivalent:*

- 1) *The principal symbol $\sigma_m(P)$ of P is invertible with respect to the convolution product for homogeneous symbols;*
- 2) *P admits a parametrix Q in $\Psi_H^{-m}(M, \mathcal{E})$, so that $PQ = QP = 1 \bmod \Psi^{-\infty}(M, \mathcal{E})$.*

Proof. First, it immediately follows from Proposition 4.9 that 2) implies 1). Conversely, in a local trivializing Heisenberg chart (4.20) shows that the invertibility of the global principal $\sigma_m(P)$ is equivalent to that of the local principal symbol. Once the latter is granted Lemma 3.7 and Proposition 3.9 allows us to carry out in a local trivializing Heisenberg chart the standard parametrix construction in a trivializing Heisenberg chart to get a parametrix for P as a Ψ_H DO of order $-m$ (see [BG, p. 142]). A classical partition of the unity argument then allows us to produce a parametrix for P in $\Psi_H^{-m}(M, \mathcal{E})$. $\square$

When a Ψ_H DO has an invertible principal symbol using the Sobolev regularity properties of its parametrices allows us to get:

Proposition 5.2 ([BG, p. 142]). *Let $P : C_c^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a Ψ_H DO of order m whose principal symbol is invertible. If $k := \Re m > 0$ then P is hypoelliptic with loss of $\frac{k}{2}$ -derivatives, i.e., for any $a \in M$, any $u \in \mathcal{E}'(M, \mathcal{E})$ and any $s \in \mathbb{R}$, we have*

$$(5.1) \quad Pu \text{ is } L_s^2 \text{ near } a \implies u \text{ is } L_{s+k}^2 \text{ near } a.$$

In particular, if M is compact then, for any reals s and s' , we have the estimate,

$$(5.2) \quad \|f\|_{L_{s+k}^2} \leq C_{ss'} (\|Pf\|_{L_s^2} + \|f\|_{L_{s'}^2}), \quad u \in C^\infty(M, \mathcal{E}).$$

Remark 5.3. We can give sharper hypoellipticity regularity results for Ψ_H DO's in terms of suitably weighted Sobolev spaces (see [FS], [Po3]). When P is a differential operator and the Levi form is non-vanishing these results correspond to the maximal hypoellipticity of P as in [HN3].

Now, let $P : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a Ψ DO of order m and assume that M is endowed with a positive density and $\mathcal{E}$ with a Hermitian metric. Let P^a be the model operator of P at a point $a \in M$ and let $\pi : G \rightarrow \mathcal{H}_\pi$ be a (nontrivial) unitary representation of $G = G_a M$. We define the symbol π_{P^a} as follows (see also [Ro1], [Gl], [CGGP]).

Let $\mathcal{H}_\pi^0(\mathcal{E}_a)$ be the subspace of $\mathcal{H}_\pi(\mathcal{E}_a) := \mathcal{H}_\pi \otimes \mathcal{E}_a$ spanned by the vectors of the form

$$(5.3) \quad \pi_f \xi = \int_G (\pi_x \otimes 1_{\mathcal{E}_a})(\xi \otimes f(x)) dx,$$

as ξ ranges over $\mathcal{H}_\pi$ and f over $\mathcal{S}_0(G, \mathcal{E}_a) = \mathcal{S}_0(G) \otimes \mathcal{E}_a$. Then we let π_{P^a} denote the (unbounded) operator of $\mathcal{H}_\pi(\mathcal{E}_a)$ with domain $\mathcal{H}_\pi^0(\mathcal{E}_a)$ such that

$$(5.4) \quad \pi_{P^a}(\pi_f \xi) = \pi_{P^a f} \xi \quad \forall f \in \mathcal{S}_0(G, \mathcal{E}_a) \quad \forall \xi \in \mathcal{H}_\pi.$$

One can check that π_{P^a*} is the adjoint of π_{P^a} on $\mathcal{H}_\pi^0$, hence is densely defined. Thus π_{P^a} is closeable and we can let $\overline{\pi_{P^a}}$ denotes its closure.

In the sequel we let $C_\pi^\infty(\mathcal{E}_a) = C_\pi^\infty \otimes \mathcal{E}_a$, where $C_\pi^\infty \subset \mathcal{H}_\pi$ denotes the space of smooth vectors of π (i.e. the subspace of vectors $\xi \in \mathcal{H}_\pi$ so that $x \rightarrow \pi(x)\xi$ is smooth from G to $\mathcal{H}_\pi$).

Proposition 5.4 ([CGGP]). 1) The domain of $\overline{\pi_{P^a}}$ always contains $C_\pi^\infty(\mathcal{E}_a)$.

2) If $\Re m \leq 0$ then the operator $\overline{\pi_{P^a}}$ is bounded.

3) We have $\overline{(\pi_{P^a})^*} = (\overline{\pi_{P^a}})^*$.

4) If P_1 and P_2 are ΨDO 's on M then $\overline{\pi_{(P_1 P_2)^a}} = \overline{\pi_{P_1^a}} \overline{\pi_{P_2^a}}$.

Remark 5.5. If $\mathcal{E}_a = \mathbb{C}$ and P^a is a differentiable operator then, as it is left-invariant, P^a belongs to the enveloping algebra $\mathcal{U}(\mathfrak{g})$ of the Lie algebra $\mathfrak{g} = \mathfrak{g}_a M$ of G . In this case $\overline{\pi_{P^a}}$ coincides on C_π^∞ with the operator $d\pi(P^a)$, where $d\pi$ is the representation of $\mathcal{U}(\mathfrak{g})$ induced by π .

Definition 5.6. We say that P satisfies the Rockland condition at a if for any nontrivial unitary irreducible representation π of $G_a M$ the operator $\overline{\pi_{P^a}}$ is injective on $C_\pi^\infty(\mathcal{E}_a)$.

Since $G = G_a M \simeq \mathbb{H}^{2n+1} \times \mathbb{R}^{d-2n}$ with $2n = \text{rk } \mathcal{L}_a$, there are left-invariant vector fields $X_0, \dots, X_d$ on G such that $X_0, \dots, X_{2n}$ are the left-invariant vector fields on $\mathbb{H}^{2n+1}$ given by (2.2) and $X_k = \frac{\partial}{\partial x_k}$ for $k = 2n+1, \dots, d$. Then, up to unitary equivalence, the nontrivial irreducible representations of G are of two types:

(i) Infinite dimensional representations $\pi^{\lambda, \xi} : G \rightarrow L^2(\mathbb{R}^n)$ parametrized by $\lambda \in \mathbb{R} \setminus 0$ and $\xi = (\xi_{2n+1}, \dots, \xi_{2n})$ such that

$$(5.5) \quad d\pi^{\lambda, \xi}(X_0) = i\lambda|\lambda|, \quad d\pi^{\lambda, \xi}(X_k) = i\lambda\xi_k, \quad k = 2n+1, \dots, d,$$

$$(5.6) \quad d\pi^{\lambda, \xi}(X_j) = |\lambda| \frac{\partial}{\partial \xi_j}, \quad d\pi^{\lambda, \xi}(X_{n+j}) = i\lambda\xi_j, \quad j = 1, \dots, n.$$

Moreover, in this case we have $C^\infty(\pi^{\pm, \xi}) = \mathcal{S}(\mathbb{R}^n)$.

(ii) One dimensional representations $\pi^\xi : G \rightarrow \mathbb{C}$ indexed by $\xi = (\xi_1, \dots, \xi_d) \in \mathbb{R}^d \setminus 0$ such that

$$(5.7) \quad d\pi^\xi(X_0) = 0, \quad d\pi^\xi(X_j) = i\xi_j, \quad j = 1, \dots, d.$$

In particular, if $P = p_m(-iX)$ with $p \in S_m(G)$ then the homogeneity of the symbol p implies that we have $\pi_P^{\lambda, \xi} = |\lambda|^m \pi_P^{\pm, \xi}$ where $\pi_P^{\pm, \xi} = \pi_P^{\pm 1, \xi}$ accordingly with the sign of λ .

On the other hand, for the representations in (ii) we have $\overline{\pi_P^\xi} = \pi_P^\xi = p_m(0, \xi_1, \dots, \xi_d)$. Therefore, we get:

Proposition 5.7. The Rockland condition for $P = p_m(-iX^a)$, $p_m \in S_m(G_a M)$, holds, and only if, the following two conditions are satisfied:

(i) The operators $\overline{\pi_P^{\pm, \xi}}$, $\xi \in \mathbb{R}^{d-2n}$, are injective on $\mathcal{S}(\mathbb{R}^n)$;

(ii) The restriction of the symbol p_m to $\{0\} \times (\mathbb{R}^n \setminus 0) \simeq H_a^* \setminus 0$ is pointwise invertible.

Remark 5.8. In the case of $G_a M = \mathbb{H}^{2n+1}$ the conditions (i) and (ii) have those considered by Taylor [Tay].

Next, if $P \in \Psi_H^m(M, \mathcal{E})$ has an invertible principal symbol, hence admits a parametrix $Q \in \Psi_H^{-m}(M, \mathcal{E})$, then for any $a \in M$ we have $Q^a P^a$ and $(Q^a)^t (P^a)^t$ are equal to 1 on $\mathcal{S}_0(G_a M, \mathcal{E}_a)$ and $\mathcal{S}_0(G_a M, \mathcal{E}_a^*)$ respectively. It then follows from Proposition 5.4 that for any nontrivial irreducible unitary representation π of G_a the operators $\overline{\pi_{P^a}}$ and $\overline{\pi_{(P^a)^t}}$ are injective on $C^\infty(\pi)$. Thus P and P^t satisfy the Rockland condition at every point of M .

Conversely, if at some point $a \in M$ the model operator $(P^a)^t$ satisfies the Rockland condition then $(P^a)^t$ is hypoelliptic (see [Ro1], [Bea], [Gl]; see also [HN1], [HN2], [CGGP]). The latter fact then implies that P^a admits a fundamental solution $k^a \in \mathcal{S}'(G_a M, \mathcal{E}_a)$ so that $P^a k^a = \delta_a$ (see [Fo], [Ge], [CGGP]). In particular, the inverse Fourier transform of k^a yields an inverse for the symbol of P^a .

This shows that if P and P^t satisfies the Rockland condition at every point then for any $a \in M$ there exists $q_{-m}^a \in S_{-m}(\mathfrak{g}_a^* M, \mathcal{E}_a)$ so that $q^a *^a \sigma_m(P)(a, \cdot) = \sigma_m(P)(a, \cdot) *^a q^a = 1$. However,

it is an open problem to determine whether q^a depends smoothly a and so define an element of $S_m(\mathfrak{g}^*M, \mathcal{E})$.

The above issue is at least true in the case of sublaplacians (see [BG] and next section). As we shall now see this result can be extended to arbitrary Ψ_H DO's when the Levi form of (M, H) has constant rank. More precisely, we have:

Proposition 5.9. *Suppose that the Levi form of (M, H) has constant rank and let $P \in \Psi_H^m(M, \mathcal{E})$. Then the following are equivalent:*

- (i) P and P^t satisfy the Rockland condition at every point of M ;
- (ii) P and P^* satisfy the Rockland condition at every point of M ;
- (iii) The principal symbol of P is invertible.

In particular, if (i) holds then P admits a parametrix in $\Psi_H^{-m}(M, \mathcal{E})$ and is hypoelliptic with loss of $\frac{1}{2}\Re m$ derivatives.

Proof. We saw that (i) implies (iii) already. The same argument shows that (ii) implies (i). Thus we only need to prove the converse statements.

Assume that P and P^t satisfies the Rockland condition at every point. For sake of simplicity let us further assume that $\mathcal{E}$ is the trivial line bundle, so that P and P^t are scalar operators. We need to show that the principal symbol of P is invertible. As this is a local issue it is enough to prove it in a local Heisenberg chart $U \subset \mathbb{R}^{d+1}$, so we may assume that P and P^t are Ψ_H DO's on U .

Moreover, since the Levi form has constant rank it follows from [Po2, Prop. 2.8] that GM is a fiber bundle of Lie group with fiber $G = \mathbb{H}^{2n+1} \times \mathbb{R}^{d-2n}$, where $2n$ is the rank of the Levi form. Therefore, by considering a trivialization of this fiber bundle by means of a suitable local H -frame (see [Po2, p. 5]) we may further assume that GU is the trivial bundle $U \times G$. In particular, the families of model operators $(P^x)_{x \in U}$ and $((P^t)^x)_{x \in U}$ can be seen as smooth families of left-invariant Ψ_H DO's on G as in [CGGP].

Now, since P^x satisfies the Rockland condition for every $x \in U$ it follows from [CGGP, Thm. 5(d)] near every $x_0 \in U$ there exists an open neighborhood $V \subset U$ of x_0 and a smooth family $(K^x)_{x \in V} \subset \mathcal{K}_{m-d-2}(G)$ such that if for $x \in V$ we let Q^x be the left-convolution operator with K^x acting on $S_0(G)$ then Q^x is a left-inverse for P^x .

As $(P^t)^x = (P^x)^t$ satisfies the Rockland condition for every $x \in V$ the same arguments show that $(P^x)^t$ is left-invertible on $S_0(G)$ for any x in an open neighborhood W of x_0 contained in V . Thus P^x is invertible with two-side inverse Q^x .

Since Q^x is the left-invariant Ψ_H DO with symbol $q_{-m}^x = (K^x)^\wedge$ in $S_{-m}(G)$ we see that we have

$$(5.8) \quad q_{-m}^x * \sigma_m(P)(x, \cdot) = \sigma_m(P)(x, \cdot) * q_{-m}^x = 1 \quad \forall x \in W.$$

In fact, as K^x depends smoothly on x , that is, yields an element of $\mathcal{K}_{-m}(W \times G)$, we get a symbol in $S_{-m}(W \times \mathfrak{g}^*)$ by letting $q_{-m}(x, \xi) = q_{-m}^x(\xi)$. Then (5.8) shows that q_{-m} is an inverse for $\sigma_m(P)$ on $W \times (\mathfrak{g}^* \setminus 0)$. This shows that $\sigma_m(P)$ is invertible near every point of U , so $\sigma_m(P)$ is an invertible symbol.

Next, since the aforementioned result of [CGGP] remains valid *mutatis standis* for systems, by working in a local trivializing Heisenberg chart we can similarly show that, in the case $\mathcal{E}$ is a general bundle, if P and P^t satisfy the Rockland condition at every point then the the principal symbol of P is invertible. Thus the assertions (i) and (ii) are equivalent.

Finally, the above arguments remain valid when we replace the transpose P^t by the adjoint of P , so the statements (ii) and (iii) are equivalent. $\square$

In particular, when the Levi form of (M, H) has constant rank and P is selfadjoint the Rockland condition for P is equivalent to the invertibility of the principal symbol of P , hence is equivalent to the existence of a parametrix in $\Psi_H^{-m}(M, \mathcal{E})$.

6. HYPOELLIPTICITY CRITERIA FOR SUBLAPLACIANS

The main focus of this section is on sublaplacians, which furnish several important examples of operators on Heisenberg manifolds. The scalar case was dealt with in [BG], but the results were not extended to sublaplacians acting on sections of vector bundles. These extensions are necessary in order to deal with sublaplacians acting on forms such as the Kohn Laplacian or the horizontal sublaplacian (see next section).

In this section, after having explained the scalar case from the point of view this paper, we extend the results to the non-scalar case. In particular, this will allow us to complete the treatment of the Kohn Laplacian in [BG] (see Remark 7.3 ahead).

Definition 6.1. *A differential operator $\Delta : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ is a sublaplacian when, near any point $a \in M$, we can put Δ in the form,*

$$(6.1) \quad \Delta = -(X_1^2 + \dots + X_d^2) - i\mu(x)X_0 + O_H(1),$$

where $X_0, X_1, \dots, X_d$ is a local H -frame of TM , the coefficient $\mu(x)$ is a local section of $\text{End } \mathcal{E}$ and the notation $O_H(1)$ means a differential operator of Heisenberg order ≤ 1 .

Let us look at the Rockland condition for a *scalar* sublaplacian $\Delta : C^\infty(M) \rightarrow C^\infty(M)$. Let $a \in M$ and let $X_0, X_1, \dots, X_d$ be a local H -frame of TM so that near a we can write

$$(6.2) \quad \Delta = -\sum_{j=1}^d X_j^2 - i\mu(x)X_0 + O_H(1),$$

where $\mu(x)$ is a smooth function near a . Using (4.10) we see that the principal symbol of Δ is

$$(6.3) \quad \sigma_2(\Delta)(x, \xi) = |\xi'|^2 + \mu(a)\xi_0, \quad \xi' = (\xi_1, \dots, \xi_d).$$

In particular we have $\sigma_2(\Delta)(x, 0, \xi') = |\xi'|^2 > 0$ for $\xi' \neq 0$, which shows that the condition (i) of Proposition 5.7 is always satisfied.

Let $L(x) = (L_{jk}(x))$ be the matrix of the Levi form $\mathcal{L}$ with respect to the H -frame $X_0, \dots, X_d$, so that for $j, k = 1, \dots, d$ we have

$$(6.4) \quad \mathcal{L}(X_j, X_k) = [X_j, X_k] = L_{jk}X_0 \mod H.$$

Equivalently, if we let $g(x)$ be the metric on H making orthonormal the frame $X_1, \dots, X_d$, then for any sections X and Y of H we have

$$(6.5) \quad \mathcal{L}(X, Y) = g(x)(L(x)X, Y)X_0 \mod H.$$

The matrix $L(x)$ is antisymmetric, so up to an orthogonal change of frame of H , which does not affect the form (6.2), we may assume that $L(a)$ is in the normal form,

$$(6.6) \quad L(a) = \begin{pmatrix} 0 & D & 0 \\ -D & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad D = \text{diag}(\lambda_1, \dots, \lambda_n), \quad \lambda_j > 0,$$

so that $\pm i\lambda_1, \dots, \pm i\lambda_n, 0, \dots, 0$ are the eigenvalues of $L(a)$ counted with multiplicity, the multiplicity of 0 being $d - 2n$. Then the model vector fields $X_0^a, \dots, X_d^a$ are:

$$(6.7) \quad X_0^a = \frac{\partial}{\partial x_0}, \quad X_k^a = \frac{\partial}{\partial x_k}, \quad k = 2n+1, \dots, d,$$

$$(6.8) \quad X_j^a = \frac{\partial}{\partial x_j} - \frac{1}{2}\lambda_j x_{n+j} \frac{\partial}{\partial x_0}, \quad X_{n+j}^a = \frac{\partial}{\partial x_j} + \frac{1}{2}\lambda_j x_j \frac{\partial}{\partial x_0}, \quad j = 1, \dots, n.$$

In terms of these vector fields the model operator of Δ at a is

$$(6.9) \quad \Delta^a = -[(X_1^a)^2 + \dots + (X_n^a)^2] - i\mu(a)X_0^a.$$

Next, under the isomorphism $\phi : \mathbb{H}^{2n+1} \times \mathbb{R}^{d-2n} \rightarrow G_a M$ given by

$$(6.10) \quad \phi(x_0, \dots, x_d) = (x_0, \lambda_1^{\frac{1}{2}} x_1, \dots, \lambda_n^{\frac{1}{2}} x_n, \lambda_1^{\frac{1}{2}} x_{n+1}, \dots, \lambda_n^{\frac{1}{2}} x_{2n}, x_{2n+1}, \dots, x_d),$$

the representations $\pi^{\pm, \xi} = \pi^{\pm 1, \xi}$, $\xi \in \{0\}^{2n} \times \mathbb{R}^{d-2n}$, become the representations of $G_a M$ such that

$$(6.11) \quad d\pi^{\pm, \xi}(X_0) = \pm i, \quad d\pi^{\pm, \xi}(X_k) = \pm i\xi_k, \quad k = 2n+1, \dots, d,$$

$$(6.12) \quad d\pi^{\pm, \xi}(X_j) = \lambda_j^{\frac{1}{2}} \frac{\partial}{\partial \xi_j}, \quad d\pi^{\pm, \xi}(X_{n+j}) = \pm i\lambda_j^{\frac{1}{2}} \xi_j, \quad j = 1, \dots, n,$$

$$(6.13) \quad \pi_{\Delta_a}^{\pm, \xi} = d\pi^{\pm, \xi}(\Delta^a) = \sum_{j=1}^n \lambda_j (-\partial_{\xi_j}^2 + \xi_j^2) \pm (\xi_{2n+1}^2 + \dots + \xi_d^2 + \mu(a)).$$

The spectrum of the harmonic oscillator $\sum_{j=1}^n \lambda_j (-\partial_{\xi_j}^2 + \xi_j^2)$ is $\sum_{j=1}^n \lambda_j (1 + 2\mathbb{N})$ and all its eigenvectors belong to $\mathcal{S}(\mathbb{R}^n)$. Thus, the operator $\pi_{\Delta_a}^{\pm, \xi}$ is injective on $\mathcal{S}(\mathbb{R}^n)$ if, and only if, $\xi_{2n+1}^2 + \dots + \xi_d^2 + \mu(a)$ is not $\pm \sum_{j=1}^n \lambda_j (1 + 2\mathbb{N})$. This occurs for any $\xi \in \{0\}^{2n} \times \mathbb{R}^{d-2n}$ if, and only if, the following condition holds

$$(6.14) \quad \mu(a) \text{ is not in the singular set } \Lambda_a,$$

$$(6.15) \quad \Lambda_a = (-\infty, -\frac{1}{2} \text{Trace } |L(a)|] \cup [\frac{1}{2} \text{Trace } |L(a)|, \infty) \quad \text{if } 2n < d,$$

$$(6.16) \quad \Lambda_a = \{\pm(\frac{1}{2} \text{Trace } |L(a)| + 2 \sum_{1 \leq j \leq n} \alpha_j |\lambda_j|); \alpha_j \in \mathbb{N}^d\} \quad \text{if } 2n = d.$$

In particular, the condition (ii) of Proposition 5.7 is equivalent to (6.14). Since the condition (i) is always satisfied, it follows that the Rockland condition for Δ is equivalent to (6.14).

Notice also that, independently of the equivalence with the Rockland condition, the condition (6.14) does not depend on the choice of the H -frame, because as Λ_a depends only on the eigenvalues of $L(a)$ which scale in the same way as $\mu(a)$ under a change of H -frame preserving the form (6.2).

On the other hand, since the transpose $(\Delta^a)^t = (\Delta^t)^a$ is given by the formula (6.9) with $\mu(a)$ replaced by $-\mu(a)$, which has no effect on (6.14), we see that the Rockland condition for $(\Delta^t)^a$ too is equivalent to (6.14). Therefore, we have obtained:

Proposition 6.2. *The Rockland conditions for Δ^t and Δ at a are both equivalent to (6.14).*

In particular, we see that if the principal symbol of Δ is invertible then the condition (6.14) holds at every point. As shown by Beals-Greiner the converse is true as well. The key result is the following.

Proposition 6.3 ([BG, Sect. 5]). *Let $U \subset \mathbb{R}^{d+1}$ be a Heisenberg chart near a and set*

$$(6.17) \quad \Omega = \{(\mu, x) \in \mathbb{C} \times U; \mu \notin \Lambda_x\}.$$

Then Ω is an open subset of $\mathbb{C} \times U$ and there exists $q_\mu(x, \xi) \in C^\infty(\Omega \times \mathbb{R}^{d+1} \setminus \{0\})$ such that:

(i) $q_\mu(x, \xi)$ is analytic with respect to μ ;

(ii) We have $q_\mu(x, \lambda\xi) = \lambda^{-2} q_\mu(x, \xi)$ for any $\lambda > 0$.

(iii) For any $(\mu, x) \in \Omega$ the symbol $q_\mu(x, \cdot)$ inverts $|\xi'|^2 + i\mu\xi_0$ on $G_x U$, i.e., we have

$$(6.18) \quad q_\mu(x, \cdot) *^x (|\xi'|^2 + i\mu\xi_0) = (|\xi'|^2 + i\mu\xi_0) *^x q_\mu(x, \cdot) = 1.$$

More precisely, $q_\mu(x, \xi)$ is obtained from the analytic continuation of the function,

$$q_\mu(x, \xi) = \int_0^\infty e^{-t\mu\xi_0} G(x, \xi, t) dt, \quad |\Re \mu| < \frac{1}{2} \operatorname{Tr} |L(x)|,$$

$$G(x, \xi, t) = \det^{-\frac{1}{2}} [\cosh(t|\xi_0||L(x)|)] \exp[-t \langle \frac{\tanh(t|\xi_0||L(x)|)}{t|\xi_0||L(x)|} \xi', \xi' \rangle].$$

This implies that if the condition (6.14) is satisfied at every point $x \in U$ then we get an inverse $q_{-2} \in S_{-2}(U \times \mathbb{R}^{d+1})$ for $\sigma_2(\Delta)(x, \xi) = |\xi'|^2 + i\mu(x)\xi_0$ on $U \times \mathbb{R}^{d+1}$ by letting

$$(6.19) \quad q_{-2}(x, \xi) = q_{\mu(x)}(x, \xi), \quad (x, \xi) \in U \times (\mathbb{R}^{d+1} \setminus 0).$$

It thus follows that if (6.14) holds at every point of M then the principal symbol of Δ is invertible near any point of M , hence admits an inverse in $S_{-2}(\mathfrak{g}^*M)$. Therefore, we get:

Proposition 6.4. *A scalar sublaplacian $\Delta : C^\infty(M) \rightarrow C^\infty(M)$ has an invertible principal symbol if, and only if, it satisfies the condition (6.14) at every point.*

Let us now extend the above results to the case of a sublaplacian $\Delta : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ acting on the sections of the vector bundle $\mathcal{E}$.

Let $a \in M$ and let $X_0, \dots, X_d$ be a local H -frame near a with respect to which we have

$$(6.20) \quad \Delta = - \sum_{j=1}^d X_j^2 - i\mu(x)X_0 + O_H(1),$$

where $\mu(x)$ is a smooth section of $\operatorname{End} \mathcal{E}$.

In a suitable basis of $\mathcal{E}_a$ the matrix of $\mu(a)$ is in triangular form,

$$(6.21) \quad \mu(a) = \begin{pmatrix} \mu_1(a) & * & * \\ 0 & \ddots & * \\ 0 & 0 & \mu_r(a) \end{pmatrix}.$$

where $\mu_1(a), \dots, \mu_r(a)$ denote the eigenvalues of $\mu(a)$ counted with multiplicity. Therefore, the model operator of Δ at a is of the form,

$$(6.22) \quad \Delta^a = \begin{pmatrix} \Delta_1^a & * & * \\ 0 & \ddots & * \\ 0 & 0 & \Delta_r^a \end{pmatrix}, \quad \Delta_j^a = -[(X_1^a)^2 + \dots + (X_d^a)^2] - i\mu_j(a)X_0^a.$$

It follows that Δ^a satisfies the Rockland condition if, and only if, so does each sublaplacian Δ_j^a , $j = 1, \dots, r$. Using Proposition 6.4 we then deduce that the Rockland condition Δ^a is equivalent to the condition,

$$(6.23) \quad \operatorname{Sp} \mu(a) \cap \Lambda_a = \emptyset.$$

Notice that the same is true for the transpose $(\Delta^a)^t$. Moreover, the condition (6.23) is independent of the choice of the basis of $\mathcal{E}_a$ or of the H -frame since the condition involves $\mu(a)$ only though its eigenvalues of $\mu(a)$ and the latter scale in the same way as that of $L(a)$ under a change of H -frame preserving the form (6.20).

Next, concerning the invertibility of the principal symbol of Δ the following extension of Proposition 6.3 holds.

Proposition 6.5. *Let $U \subset \mathbb{R}^{d+1}$ be a trivializing Heisenberg chart near a and set*

$$(6.24) \quad \Omega = \{(\mu, x) \in M_r(\mathbb{C}) \times U; \mu \notin \Lambda_x\}.$$

Then Ω is an open subset of $M_r(\mathbb{C}) \times U$ and there exists $q_\mu(x, \xi) \in C^\infty(\Omega \times \mathbb{R}^{d+1} \setminus 0, M_r(\mathbb{C}))$ so that:

(i) $q_\mu(x, \xi)$ is analytic with respect to μ ;

(ii) We have $q_\mu(x, \lambda\xi) = \lambda^{-2}q_\mu(x, \xi)$ for any $\lambda > 0$.

(iii) For any $(\mu, x) \in \Omega$ the symbol $q_\mu(x, \cdot)$ inverts $|\xi'|^2 + i\mu\xi_0$ on $G_x U$, that is, we have

$$(6.25) \quad q_\mu(x, \cdot) *^x (|\xi'|^2 + i\mu\xi_0) = (|\xi'|^2 + i\mu\xi_0) *^x q_\mu(x, \cdot) = 1.$$

Proof. It is enough to prove that near point $(\mu_0, x_0) \in \Omega$ there exists an open neighborhood Ω' contained in Ω and a function $q_\mu(x, \xi) \in C^\infty(\Omega \times \mathbb{R}^{d+1} \setminus 0, M_r(\mathbb{C}))$ satisfying the properties (i), (ii) and (iii) on $\Omega' \times \mathbb{R}^{d+1} \setminus 0$.

To this end observe that since $\text{Sp } \mu \subset \overline{B}(0, \|\mu\|)$ for any $\mu \in M_r(\mathbb{C})$, we see that if we let $K = \overline{B}(0, \|\mu_0\| + 1)$ then any $\mu \in M_r(\mathbb{C})$ close enough to μ_0 has its spectrum contained in K .

Let $\delta > 0$ be small enough so that $\delta < \frac{1}{2} \text{dist}(\text{Sp } \mu_0, \Lambda_{x_0})$ and define $V_1 = \text{Sp } \mu_0 + D(0, \delta)$ and $V_2 = \Lambda_{x_0} + D(0, \delta)$, so that V_1 and V_2 are disjoint open subsets of $\mathbb{C}$ containing $\text{Sp } \mu_0$ and Λ_{x_0} respectively.

Notice that for any μ close enough to μ_0 we have $\text{Sp } \mu \subset V_1$. Otherwise there exists a sequence $(\mu_k)_{k \geq 1} \subset M_r(\mathbb{C})$ converging to μ_0 and a sequence of eigenvalues $(\lambda_k)_{k \geq 1} \subset K$, $\lambda_k \in \text{Sp } \mu_k$, such that $\lambda_k \notin V_1$ for any $k \geq 1$. Since the sequence $(\lambda_k)_{k \geq 1}$ is bounded, we may assume that it converges to some $\lambda \notin V_1$. Necessarily λ is an eigenvalue of μ_0 , which contradicts the fact that $\lambda \notin V_1$. Thus there exists $\eta_1 > 0$ so that for any $\mu \in B(\mu_0, \eta_1)$ we have $\text{Sp } \mu \subset V_1$.

Similarly there exists $\eta_2 > 0$ so that for any $x \in B(x_0, \eta_2)$ we have $\text{Sp } |L(x)| \subset \text{Sp } |L(x_0)| + D(0, \delta)$, which implies $\Lambda_x \subset \Lambda_{x_0} + D(0, \delta) = V_2$. Therefore the open set $\Omega' = B(\mu_0, \eta_1) \times B(x_0, \eta_2)$ is such that for any $(\mu, x) \in \Omega'$ we have $\text{Sp } \mu \cap \Lambda_x \subset V_1 \cap V_2 = \emptyset$, that is, Ω' is an open neighborhood of (μ_0, x_0) contained in Ω .

Next, let Γ be a smooth curve of index 1 such that the bounded connected component of $\mathbb{C} \setminus \Gamma$ contains V_1 and its unbounded component contains V_2 . Then we define an element of $\text{Hol}(B(\mu_0, \eta_1)) \hat{\otimes} C^\infty(B(x_0, \eta_2) \times \mathbb{R}^{d+1} \setminus 0)$ by letting

$$(6.26) \quad q_\mu(x, \xi) = \frac{1}{2i\pi} \int_\Gamma q_\gamma(x, \xi)(\gamma - \mu) d\gamma, \quad (\mu, x, \xi) \in \Omega' \times \mathbb{R}^{d+1} \setminus 0.$$

This function is homogeneous of degree -2 with respect to ξ and for any $(\mu, x) \in \Omega'$ we have

$$(6.27) \quad q_\mu(x, \cdot) *^x (|\xi'|^2 + i\mu\xi_0) = \frac{1}{2i\pi} \int_\Gamma q_\gamma(x, \cdot) *^x (|\xi'|^2 + i\mu\xi_0)(\gamma - \mu)^{-1} d\gamma, \\ \frac{1}{2i\pi} \int_\Gamma [(\gamma - \mu)^{-1} - iq_\gamma(x, \cdot) * \xi_0] d\gamma = 1.$$

Similarly we have $(|\xi'|^2 + i\mu\xi_0) *^x q_\mu(x, \cdot) = 1$. Thus $q_\mu(x, \xi)$ satisfies the properties (i), (ii) and (iii) on $\Omega' \times \mathbb{R}^{d+1} \setminus 0$. The proof is thus complete. $\square$

In the same way as Proposition 6.3 in the scalar case, Proposition 6.5 implies that when the condition (6.23) holds everywhere the principal symbol of Δ admits an inverse in $S_{-2}(\mathfrak{g}^*M, \mathcal{E})$. We have thus proved:

Proposition 6.6. 1) At every point $a \in M$ the Rockland conditions for Δ and Δ^t are equivalent to (6.23).

2) The principal symbol of Δ is invertible if, and only if, the condition (6.23) holds everywhere. Moreover, when the latter occurs Δ admits a parametrix in $\Psi_H^{-2}(M, \mathcal{E})$ and is hypoelliptic with loss of 1 derivative.

7. EXAMPLES OF HYPOELLIPTIC OPERATORS ON HEISENBERG MANIFOLDS

In this section we explain how the previous results of this paper can be used to deal with the hypoellipticity for the main geometric operators on Heisenberg manifolds: Hörmander's sum of

squares, Kohn Laplacian, horizontal sublaplacian and contact Laplacian. In particular, the treatment in [BG] of the Kohn Laplacian and we establish a criterion for the invertibility of the horizontal sublaplacian, which has not been done before.

7.1. Hörmander's sum of squares. Let $X_1, \dots, X_m$ be (real) vector fields on a manifold M^{d+1} and consider the sum of squares,

$$(7.1) \quad \Delta = -(X_1^2 + \dots + X_m^2).$$

By a celebrated theorem of Hörmander [Hö2] the operator Δ is hypoelliptic provided that the following bracket condition is satisfied: the vector fields $X_0, X_1, \dots, X_m$ together with their successive Lie brackets $[X_{j_1}, [X_{j_2}, \dots, X_{j_l}] \dots]$ span the tangent bundle TM at every point.

When $X_1, \dots, X_m$ span a hyperplane bundle H the operator Δ is a sublaplacian with *real* coefficients and the bracket condition reduces to $H + [H, H] = TM$, which is equivalent to the non-vanishing of the Levi form of (M, H) .

More generally, given a vector bundle $\mathcal{E}$, the theorem of Hörmander holds for sublaplacians $\Delta : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ of the form

$$(7.2) \quad \Delta = -(\nabla_{X_1}^2 + \dots + \nabla_{X_m}^2) + O_H(1),$$

where ∇ is a connection on $\mathcal{E}$. In particular, if M is endowed with a positive density and $\mathcal{E}$ with a Hermitian metric, this includes the selfadjoint sum of squares,

$$(7.3) \quad \Delta = \nabla_{X_1}^* \nabla_{X_1} + \dots + \nabla_{X_m}^* \nabla_{X_m}.$$

In fact, if Δ is a sublaplacian of the form (7.2) then in (6.1) the matrix $\mu(x)$ vanishes, so that (6.23) holds if, and only if, the Levi form does not vanish at x . Therefore, we obtain:

Proposition 7.1. *Let $\Delta : C^\infty(M, \mathcal{E}) \rightarrow C^\infty(M, \mathcal{E})$ be a (generalized) sum of squares of the form (7.2). Then:*

1) *At a point $x \in M$ the operators Δ and Δ^t satisfies the Rockland condition if, and only if, the Levi form $\mathcal{L}$ does not vanish at x .*

2) *The principal symbol of Δ is invertible if, and only if, the Levi form is non-vanishing. In particular, when the latter occurs Δ admits a parametrix in $\Psi_H^{-2}(M, \mathcal{E})$ and is hypoelliptic with loss of one derivative.*

In particular, since the nonvanishing of the Levi form is equivalent to the bracket condition $H + [H, H] = TM$, we see that, the special case of Heisenberg manifolds, we recover the hypoellipticity result of [Hö2] for sums of squares.

7.2. The Kohn Laplacian. In [KR] Kohn-Rossi showed that the Dolbeault complex on a bounded complex domain induces on its boundary a horizontal complex of differential forms. This was later extended by Kohn [Koh1] to the general setting of a CR manifold M^{2n+1} as follows.

Let M^{2n+1} be a CR manifold with CR bundle $T_{1,0} \subset T_{\mathbb{C}}M$ and set $T_{0,1} = \overline{T_{1,0}}$. Then the subbundle $H = \Re(T_{1,0} \oplus T_{0,1}) \subset TM$ admits an integrable complex structure and the splitting $H \otimes \mathbb{C} = T_{1,0} \oplus T_{0,1}$ gives rise to a decomposition $\Lambda H^* \otimes \mathbb{C} = \oplus_{0 \leq p, q \leq n} \Lambda^{p,q}$, where (p, q) is called a bidegree of a form with values in $\Lambda^{p,q}$.

Assume that $T_{\mathbb{C}}M$ is endowed with a Hermitian metric such that $T_{1,0}$ and $T_{0,1}$ are orthogonal subspaces and complex conjugation is an (antilinear) isometry. This Hermitian metric gives rise to a Hermitian metric on $\Lambda^* T_{\mathbb{C}}^* M$ with respect to which the decomposition $\Lambda H^* \otimes \mathbb{C} = \oplus_{0 \leq p, q \leq n} \Lambda^{p,q}$ becomes orthogonal. Let $\Pi_{p,q} : \Lambda^* T_{\mathbb{C}}^* M \rightarrow \Lambda^{p,q}$ be the orthogonal projection onto $\Lambda^{p,q}$. Then the Kohn-Rossi operator $\bar{\partial}_b : C^\infty(M, \Lambda^{p,q}) \rightarrow C^\infty(M, \Lambda^{p,q+1})$ is given by

$$(7.4) \quad \bar{\partial}_b \eta = \Pi_{p,q+1}(d\eta), \quad \eta \in C^\infty(M, \Lambda^{p,q}).$$

Since the integrability of $T_{1,0}$ implies that $\bar{\partial}_b^2 = 0$, this yields chain complexes $\bar{\partial}_b : C^\infty(M, \Lambda^{p,*}) \rightarrow C^\infty(M, \Lambda^{p,*+1})$.

Endowing M with a smooth density $\rho > 0$ we let $\bar{\partial}_b^*$ denote the formal adjoint of $\bar{\partial}_b$. Then the Kohn Laplacian is

$$(7.5) \quad \square_b = (\bar{\partial}_b + \bar{\partial}_b^*)^2 = \bar{\partial}_b^* \bar{\partial}_b + \bar{\partial}_b \bar{\partial}_b^*.$$

The Kohn Laplacian is a sublaplacian (see [FS, Sect. 13], [BG, Sect. 20]) and it was shown by Kohn [Koh1] that under the condition $Y(q)$ this operator is hypoelliptic with loss of one derivative when acting on (p, q) -forms.

The condition $Y(q)$ means that if for $x \in M$ we let $(r(x) - \kappa(x), \kappa(x), n - r(x))$ be the signature of L_θ at x , so that $r(x)$ is the rank of L_θ and $\kappa(x)$ the number of its negative eigenvalues, then the condition $Y(q)$ is satisfied at $x \in M$ when we have

$$(7.6) \quad q \notin \{\kappa(x), \kappa(x) + 1, \dots, \kappa(x) + n - r(x)\} \cup \{r(x) - \kappa(x), r(x) - \kappa(x) + 1, \dots, n - \kappa(x)\}.$$

For instance, when M is κ -strictly pseudoconvex, the $Y(q)$ -condition exactly means that we must have $q \neq \kappa$ and $q \neq n - \kappa$.

In fact, as shown in [BG, Sect. 21], at every point $a \in M$ the condition $Y(q)$ is equivalent to the condition (6.23) for the Kohn Laplacian acting on (p, q) -forms. Therefore, using Proposition 6.6 we immediately get:

Proposition 7.2. *Let $\square_b : C^\infty(M, \Lambda^{p,q}) \rightarrow C^\infty(M, \Lambda^{p,q})$ be the Kohn Laplacian acting on (p, q) -forms.*

- 1) *At a point $x \in M$ the Rockland condition for $\square_b$ is equivalent to the condition $Y(q)$.*
- 2) *The principal symbol of $\square_b$ is invertible if, and only if, the condition $Y(q)$ is satisfied at every point. In particular, when the latter occurs $\square_b$ admits a parametrix in $\Psi_H^{-2}(M, \Lambda^{p,q})$ and is hypoelliptic with loss of one derivative.*

Remark 7.3. The proof of the second part of the statement in [BG, Sect. 21] is not quite complete, because Beals-Greiner claim that diagonalizing the leading part of the Kohn Laplacian allows us to use to apply the results for scalar sublaplacians. While this property is true in case of a Levi Metric (see [FS]), it may fail for a general metric on $T_{\mathbb{C}}M$ since a smooth eigenframe needs not exists. For instance, for the Kohn Laplacian the eigenvalues of the matrix $\mu(x)$ in (6.1) with respect to an orthonormal H -frame of TM are given in terms of eigenvalues of the Levi form (see Eq. (21.31) in [BG]), but the latter need not depend smoothly on x (unless the metric on $T_{\mathbb{C}}M$ is a Levi metric). Therefore, in order to deal with the Kohn Laplacian acting on forms, we really need to use the version for sublaplacians acting on section of vector bundles, provided by Proposition 6.6, but not deal with in [BG].

Remark 7.4. The $Y(q)$ -condition is only a sufficient condition for the hypoellipticity of $\square_b$, as the latter may be hypoelliptic when the $Y(q)$ -condition fails (e.g. [Koh2], [Ko], [Ni]).

7.3. The horizontal sublaplacian on a Heisenberg manifold. Let (M^{d+1}, H) be a Heisenberg manifold endowed with a Riemannian metric and let $\Lambda_{\mathbb{C}}^* H^* = \bigoplus_{k=0}^d \Lambda_{\mathbb{C}}^k H^*$ be the (complexified) bundle of horizontal forms. Then the horizontal sublaplacian $\Delta_b : C^\infty(M, \Lambda_{\mathbb{C}}^* H^*) \rightarrow C^\infty(M, \Lambda_{\mathbb{C}}^* H^*)$ is given by

$$(7.7) \quad \Delta_b = d_b^* d_b + d_b d_b^*, \quad d_b \alpha = \pi_b(d\alpha),$$

where π_b denotes the orthogonal projection of $\Lambda_{\mathbb{C}}^* T^* M$ onto $\Lambda_{\mathbb{C}}^* H^*$.

This operator was first introduced by Tanaka [Ta] for strictly pseudoconvex CR manifolds, but versions of this operator acting on functions were independently defined by Greenleaf [Gr] and Lee [Le]. Moreover, it can be shown that $d_b^2 = 0$ if, and only if, the subbundle H is integrable, so in general Δ_b is not the Laplacian of a chain complex.

As we shall see to determine under which condition the principal symbol of Δ_b is invertible the relevant condition to look at is the $X(k)$ -condition below.

Definition 7.5. For $x \in M$ let $2r(x)$ denote the rank of the Levi form $\mathcal{L}$ at x . Then we say that $\mathcal{L}$ satisfies the condition $X(k)$ at x when we have

$$(7.8) \quad k \notin \{r(x), r(x) + 1, \dots, d - r(x)\}.$$

For instance, the condition $X(0)$ is satisfied if, and only if, the Levi form does not vanish. Also, if M^{2n+1} is a contact manifold or a nondegenerate CR manifold then the Levi form is everywhere nondegenerate, so $r(x) = 2n$ and the $X(k)$ -condition becomes $k \neq n$. In any case, we have:

Proposition 7.6. Let $\Delta_b : C^\infty(M, \Lambda_{\mathbb{C}}^k H^*) \rightarrow C^\infty(M, \Lambda_{\mathbb{C}}^k H^*)$ be the horizontal sublaplacian acting on horizontal forms of degree k .

1) At a point $x \in M$ the Rockland condition for Δ_b is equivalent to the condition $X(k)$.

2) The principal symbol of Δ_b is invertible if, and only if, the condition $X(k)$ is satisfied at every point. In particular, when the latter occurs Δ_b admits a parametrix in $\Psi_H^{-2}(M, \Lambda_{\mathbb{C}}^k H^*)$ and is hypoelliptic with loss of one derivative.

Proof. First, thanks to Proposition 6.6 we only have to check that for $k = 0, \dots, d$ at any point a the condition (6.23) for $\Delta_b|_{\Lambda_{\mathbb{C}}^k H^*}$ is equivalent to the condition $X(k)$.

Next, let $U \subset \mathbb{R}^{d+1}$ be a Heisenberg chart around a together with an orthonormal H -frame $X_0, X_1, \dots, X_d$ of TU . Let g be the Riemannian metric of M . Then on U we can write the Levi form $\mathcal{L}$ in the form,

$$(7.9) \quad \mathcal{L}(X, Y) = [X, Y] = \langle L(x)X, Y \rangle X_0 \mod H,$$

for some antisymmetric section $L(x)$ of $\text{End}_{\mathbb{R}} H$. In particular, if for $j, k = 1, \dots, d$ we let $L_{jk} = \langle LX_j, X_k \rangle$ then we have

$$(7.10) \quad [X_j, X_k] = L_{jk}X_0 \mod H.$$

Let $2n$ be the rank of $L(a)$. Since the condition (6.23) for Δ_b at a is independent of the choice of the Heisenberg chart, we may assume that U is chosen in such way that at $x = a$ we have $g(a) = 1$ and $L(a)$ is in the normal form,

$$(7.11) \quad L(a) = \begin{pmatrix} 0 & D & 0 \\ -D & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad D = \text{diag}(\lambda_1, \dots, \lambda_n), \quad \lambda_j > 0,$$

so that $\pm i\lambda_1, \dots, \pm i\lambda_n$ are the nonzero eigenvalues of $L(a)$ counted with multiplicity.

Let $\omega^1, \dots, \omega^n$ be the coframe of H^* dual to $X_1, \dots, X_d$. For a 1-form ω we let $\varepsilon(\omega)$ denote the exterior product and $\iota(\omega)$ denote the interior product with ω , that is, the contraction with the vector fields dual to ω . For an ordered subset $J = \{j_1, \dots, j_k\} \subset \{1, \dots, d\}$, so that $j_1 < \dots < j_d$, we let $\omega^J = \omega^{j_1} \wedge \dots \wedge \omega^{j_k}$ (we make the convention that $\omega^\emptyset = 1$). Then the forms ω^J 's give rise to an orthonormal frame of $\Lambda_{\mathbb{C}}^* H^*$ over U . With respect to this frame we have

$$(7.12) \quad d_b = \sum_{j=1}^d \varepsilon(\omega^j) X_j \quad \text{and} \quad d_b = - \sum_{l=1}^d \iota(\omega^l) X_l + O_H(1).$$

Thus,

$$(7.13) \quad \Delta_b = d_b^* d_b + d_b d_b^* = - \sum_{j,l} [\varepsilon(\omega^j) \iota(\omega^l) X_j X_l + \iota(\omega^l) \varepsilon(\omega^j) X_l X_j] + O_H(1) =$$

$$- \frac{1}{2} \sum_{j,l=1}^d [(\varepsilon(\omega^j) \iota(\omega^l) + \iota(\omega^l) \varepsilon(\omega^j))(X_j X_l + X_l X_j) + (\varepsilon(\omega^j) \iota(\omega^l) - \iota(\omega^l) \varepsilon(\omega^j))[X_j, X_l]] + O_H(1).$$

Combining this with (7.10) and the relations,

$$(7.14) \quad \varepsilon(\omega^j)\iota(\omega^l) + \iota(\omega^l)\varepsilon(\omega^j) = \delta_{jl}, \quad j, l = 1, \dots, d,$$

we then obtain

$$(7.15) \quad \Delta_b = - \sum_{j=1}^d X_j^2 - i\mu(x)X_0 + O_H(1), \quad \mu(x) = \frac{1}{i} \sum_{j,l=1}^d \varepsilon(\omega^j)\iota(\omega^l)L_{jl}.$$

In particular, thanks to (7.11) at $x = a$ we have

$$(7.16) \quad \mu(a) = \frac{1}{i} \sum_{j=1}^n (\varepsilon(\omega^j)\iota(\omega^{n+j}) - \varepsilon(\omega^{n+j})\iota(\omega^j))\lambda_j.$$

For $j = 1, \dots, n$ let $\theta^j = \frac{1}{\sqrt{2}}(\omega^j + i\omega^{n+j})$ and $\bar{\theta}^j = \frac{1}{\sqrt{2}}(\omega^j - i\omega^{n+j})$. Then we have:

$$(7.17) \quad \begin{aligned} & \frac{1}{i}(\varepsilon(\omega^j)\iota(\omega^{n+j}) - \varepsilon(\omega^{n+j})\iota(\omega^j)) = \\ & \frac{-1}{2}[(\varepsilon(\theta^j) + \varepsilon(\bar{\theta}^j))(\iota(\bar{\theta}^j) - \iota(\theta^j)) - (\varepsilon(\theta^j) - \varepsilon(\bar{\theta}^j))(\iota(\bar{\theta}^j) + \iota(\theta^j))] = \varepsilon(\theta^j)\iota(\theta^j) - \varepsilon(\bar{\theta}^j)\iota(\bar{\theta}^j). \end{aligned}$$

Thus,

$$(7.18) \quad \mu(a) = \sum_{j=1}^n (\varepsilon(\theta^j)\iota(\theta^j) - \varepsilon(\bar{\theta}^j)\iota(\bar{\theta}^j))\lambda_j.$$

For any ordered subset $J = \{j_1, \dots, j_p\}$ of $\{1, \dots, n\}$ we let

$$(7.19) \quad \theta^J = \theta^{j_1} \wedge \dots \wedge \theta^{j_p}, \quad \theta^{\bar{K}} = \theta^{\bar{k}_1} \wedge \dots \wedge \theta^{\bar{k}_q},$$

Then the forms $\theta^J \wedge \theta^{\bar{K}} \wedge \omega^L$ give rise to an orthonormal frame of $\Lambda_{\mathbb{C}}^* H^*$ as J and K range over all the ordered subsets of $\{1, \dots, n\}$ and L over all the ordered subsets of $\{2n+1, \dots, d\}$. For $j = 1, \dots, n$ we have

$$(7.20) \quad \varepsilon(\theta^j)\iota(\theta^j)(\theta^J \wedge \theta^{\bar{K}} \wedge \omega^L) = \begin{cases} \theta^J \wedge \theta^{\bar{K}} \wedge \omega^L & \text{if } j \in J, \\ 0 & \text{if } j \notin J, \end{cases}$$

$$(7.21) \quad \varepsilon(\bar{\theta}^j)\iota(\bar{\theta}^j)(\theta^J \wedge \theta^{\bar{K}} \wedge \omega^L) = \begin{cases} \theta^J \wedge \theta^{\bar{K}} \wedge \omega^L & \text{if } j \in K, \\ 0 & \text{if } j \notin K. \end{cases}$$

Combining this with (7.18) then gives

$$(7.22) \quad \mu(a)(\theta^J \wedge \theta^{\bar{K}} \wedge \omega^L) = \mu_{J,\bar{K}}(a)\theta^J \wedge \theta^{\bar{K}} \wedge \omega^L, \quad \mu_{J,\bar{K}}(a) = \sum_{j \in J} \lambda_j - \sum_{j \in K} \lambda_j.$$

This shows that $\mu(a)$ diagonalizes in the basis of $\Lambda_{\mathbb{C}}^* H_a^*$ provided by the forms of $\theta^J \wedge \theta^{\bar{K}} \wedge \omega^L$ with eigenvalues given by the numbers $\mu_{J,\bar{K}}(a)$. In particular, for $k = 0, \dots, d$ we have

$$(7.23) \quad \text{Sp } \mu(a)|_{\Lambda^k H^*} = \{\mu_{J,\bar{K}}; |J| + |K| \leq k\}.$$

Note that we always have $|\mu_{J,K}| \leq \sum_{j=1}^n \lambda_j$ with equality if, and only if, one the subsets J or K is empty and the other is $\{1, \dots, n\}$, which occurs for eigenvectors in the subspace spanned by the forms $\theta^1 \wedge \dots \wedge \theta^n \wedge \omega^L$ and $\bar{\theta}^1 \wedge \dots \wedge \bar{\theta}^n \wedge \omega^L$ as L ranges over all the subsets of $\{2n+1, \dots, d\}$.

Since $\lambda_1, \dots, \lambda_n$ are the eigenvalues of $|L(a)|$, each of them counted twice, it follows that the condition (6.23) for $\Delta_b|_{\Lambda_{\mathbb{C}}^k H^*}$ reduces to $\pm \sum_{j=1}^n \lambda_j \notin \text{Sp } \mu(a)|_{\Lambda^k H^*}$. This latter condition is satisfied if, and only if, the space $\Lambda_{\mathbb{C}}^k H_a^*$ does contain any of the forms $\theta^1 \wedge \dots \wedge \theta^n \wedge \omega^L$ and $\bar{\theta}^1 \wedge \dots \wedge \bar{\theta}^n \wedge \omega^L$ with L subset of $\{2n+1, \dots, d\}$. Therefore, the sublaplacian $\Delta_b|_{\Lambda_{\mathbb{C}}^k H^*}$ satisfies (6.23) at a if, and

only if, the integer k is not between n and $n + d - 2n = d - n$, that is, if, and only if, the condition $X(k)$ holds at a . The proof is thus achieved. $\square$

Finally, suppose that M is a CR manifold with Heisenberg structure $H = \Re(T_{1,0} \oplus T_{0,1})$ and assume that $T_{\mathbb{C}}M$ is endowed with a Hermitian metric with respect to which $T_{1,0}$ and $T_{0,1}$ are orthogonal subspaces and complex conjugation is an isometry. Then we have $d_b = \bar{\partial}_b + \partial_b$, where ∂_b denotes the conjugate of $\bar{\partial}_b$, that is, $\partial_b \alpha = \overline{\bar{\partial}_b \bar{\omega}}$ for any $\omega \in C^\infty(M, \Lambda_{\mathbb{C}}^* H^*)$. Moreover, as $\bar{\partial}_b \partial_b^* + \partial_b^* \bar{\partial}_b = \bar{\partial}_b^* \partial_b + \partial_b \bar{\partial}_b^* = 0$ (see [Ta]) we get

$$(7.24) \quad \Delta_b = \square_b + \bar{\square}_b,$$

where $\bar{\square}_b$ is the conjugate of $\square_b$. In particular, we see that the horizontal sublaplacian preserves the bidegree, i.e., it acts on (p, q) -forms.

In fact, along the similar lines as that of [BG, pp. 151–156] and of the proof of Proposition 7.6 above, one can show that the condition (6.23) at a point $x \in M$ for Δ_b acting on (p, q) -forms is equivalent to the condition $Y(p, q)$ below:

$$(7.25) \quad \{(p, q), (q, p)\} \cap \{(\kappa(x) + j, r(x) - \kappa(x) + k); \max(j, k) \leq n - r(x)\} = \emptyset,$$

where $(r(x) - \kappa(x), \kappa(x), n - r(x))$ is the signature at x of the Levi form L_θ associated to some non-vanishing real 1-form θ annihilating $T_{1,0} \oplus T_{0,1}$. In particular, when M is κ -strictly pseudoconvex the $Y(p, q)$ reduces to $(p, q) \neq (\kappa, n - \kappa)$ and $(p, q) \neq (n - \kappa, \kappa)$.

7.4. Contact complex and the contact Laplacian. Given an orientable contact manifold (M^{2n+1}, θ) the contact complex of Rumin [Ru] can be seen as an attempt to get on M a complex of horizontal differential forms by forcing out the equalities $d_b^2 = 0$ and $(d_b^*)^2 = 0$ as follows.

Let $H = \ker \theta$ and assume that H is endowed with a calibrated almost complex structure $J \in \text{End}_{\mathbb{R}} H$, $J^2 = -1$, so that $d\theta(X, JX) = -d\theta(JX, X) > 0$ for any section X of H . We then can endow M with the Riemannian metric $g_\theta = d\theta(\cdot, J\cdot) + \theta^2$.

In addition, let T be the Reeb fields of θ . Then we have

$$(7.26) \quad d_b^2 = -\mathcal{L}_T \varepsilon(d\theta) = \varepsilon(d\theta) \mathcal{L}_T,$$

where $\varepsilon(d\theta)$ denotes the exterior multiplication by $d\theta$.

There are two ways of modifying the space $\Lambda_{\mathbb{C}}^* H^*$ of horizontal forms to get a complex. The first one is to force the equality $d_b^2 = 0$ by restricting the operator d_b to $\Lambda_2^* := \ker \varepsilon(d\theta) \cap \Lambda_{\mathbb{C}}^* H^*$ since this bundle is stable under d_b and on there d_b^2 vanishes.

The second way is to similarly force the equality $(d_b^*)^2 = 0$ by restricting d_b^* to $\Lambda_1^* := \ker \iota(d\theta) \cap \Lambda_{\mathbb{C}}^* H^* = (\text{im } \varepsilon(d\theta))^\perp \cap \Lambda_{\mathbb{C}}^* H^*$, where $\iota(d\theta)$ denotes the interior product with $d\theta$. This amounts to replace d_b by the operator $\pi_1 \circ d_b$, where π_1 is the orthogonal projection onto Λ_1^* .

On the other hand, since $d\theta$ is nondegenerate on H the operator $\varepsilon(d\theta) : \Lambda_{\mathbb{C}}^k H^* \rightarrow \Lambda_{\mathbb{C}}^{k+2} H^*$ is injective for $k \leq n - 1$ and surjective for $k \geq n + 1$. This implies that $\Lambda_2^k = 0$ for $k \leq n$ and $\Lambda_1^k = 0$ for $k \geq n + 1$. Therefore, we only have two halves of complexes. The key observation of Rumin is that we get a full complex by connecting the two halves by means of the (second order) differential operator,

$$(7.27) \quad D_R : \Lambda_1^n \longrightarrow \Lambda_2^{n+1}, \quad D_R = \mathcal{L}_T - d_b \varepsilon(d\theta)^{-1} d_b,$$

where $\varepsilon(d\theta)^{-1}$ is the inverse of $\varepsilon(d\theta) : \Lambda_{\mathbb{C}}^{n-1} H^* \rightarrow \Lambda_{\mathbb{C}}^{n+1} H^*$. In other words, letting $\Lambda^k = \Lambda_1^k$ for $k = 0, \dots, n - 1$ and $\Lambda^k = \Lambda_2^k$ for $k = n + 1, \dots, 2n$, we have the chain complex,

$$(7.28) \quad C^\infty(M) \xrightarrow{d_R} \dots \xrightarrow{d_R} C^\infty(M, \Lambda_1^n) \xrightarrow{D_R} C^\infty(M, \Lambda_2^n) \xrightarrow{d_R} \dots \xrightarrow{d_R} C^\infty(M, \Lambda^{2n}),$$

where $d_R : C^\infty(M, \Lambda^k) \rightarrow C^\infty(M, \Lambda^{k+1})$ is equal to $\pi_1 \circ d_b$ for $k = 0, \dots, n - 1$ and to d_b for $k = n + 1, \dots, 2n$. This complex is called the contact complex of M .

The contact Laplacian $\Delta_R : C^\infty(M, \Lambda^* \oplus \Lambda_*^n) \rightarrow C^\infty(M, \Lambda^* \oplus \Lambda_*^n)$ is given by the formulas,

$$(7.29) \quad \Delta_R = \begin{cases} (n-k)d_R d_R^* + (n-k+1)d_R^* d_R & \text{on } \Lambda^k, k = 0, \dots, n-1, \\ (d_R^* d_R)^2 + D_R^* D_R & \text{on } \Lambda_1^n, \\ D_R D_R^* + (d_R d_R^*) & \text{on } \Lambda_2^n, \\ (n-k+1)d_R d_R^* + (n-k)d_R^* d_R & \text{on } \Lambda^k, k = n+1, \dots, 2n. \end{cases}$$

By comparing the contact Laplacian Δ_R to the horizontal sublaplacian Δ_b Rumin [Ru] was able to show that on every degree Δ_R satisfies the Rockland condition at every point. He then used results of Helffer-Nourrigat ([HN3]) to show that Δ_R was maximal hypoelliptic.

Alternatively, since in the contact case both $d\theta$ and $\mathcal{L}$ are nondegenerate on H , once the Rockland condition is granted we may directly apply Proposition 5.9 to get:

Proposition 7.7. 1) For $k = 0, \dots, 2n$ with $k \neq n$ the contact Laplacian Δ_R acting on sections of Λ^k has an invertible principal symbol of degree -2 , hence admits a parametrix in $\Psi_H^{-2}(M, \Lambda^k)$ and is hypoelliptic with loss of one derivative.

2) For $k = n$ the contact Laplacian Δ_R acting on sections of Λ_*^n has an invertible principal symbol of degree -4 , hence admits a parametrix in $\Psi_H^{-4}(M, \Lambda_*^n)$ and is hypoelliptic with loss of two derivatives.

8. PROOF OF PROPOSITION 3.18

First, we need the lemma below.

Lemma 8.1. For $\Re m > 0$ we have $\mathcal{K}^m(U \times \mathbb{R}^{d+1}) \subset C^\infty(U) \hat{\otimes} C^{[\frac{\Re m}{2}]}(\mathbb{R}^{d+1})$.

Proof. Let $N = [\frac{\Re m}{2}]$ and let α be a multi-order such that $|\alpha| \leq N$. As $\langle \alpha \rangle \leq 2|\alpha| \leq -(k+d+2)$ the multiplication by ξ^α maps $S^{\hat{m}}(U \times \mathbb{R}^{d+1})$ to $C^\infty(U) \hat{\otimes} L^1(\mathbb{R}^{d+1})$. Composing it with the inverse Fourier transform with respect to ξ then shows that the map $p \rightarrow \partial_y^\alpha \check{p}_{\xi \rightarrow y}$ maps $S^{\hat{m}}(U \times \mathbb{R}^{d+1})$ to $C^\infty(U) \hat{\otimes} C^0(\mathbb{R}^{d+1})$. It then follows that for any $p \in S^{\hat{m}}(U \times \mathbb{R}^{d+1})$ the transform $\check{p}_{\xi \rightarrow y}(x, y)$ belongs to $C^\infty(U) \hat{\otimes} C^N(\mathbb{R}^{d+1})$.

Now, if $K \in \mathcal{K}^m(U \times \mathbb{R}^{d+1})$ then by Lemma 3.14 there exists $p \in S^{\hat{m}}(U \times \mathbb{R}^{d+1})$, $\hat{m} = -(m+d+2)$, such that $K(x, y)$ is equal to $\check{p}_{\xi \rightarrow y}(x, y)$ modulo a smooth function. Hence $K(x, y)$ belongs to $C^\infty(U) \hat{\otimes} C^N(\mathbb{R}^{d+1})$. The lemma is thus proved. $\square$

We are now ready to prove Proposition 3.18. Let $\tilde{U}$ be an open subset of $\mathbb{R}^{d+1}$ together with a hyperplane bundle $\tilde{H} \subset T\tilde{U}$ and a $\tilde{H}$ -frame of $T\tilde{U}$ and let $\phi : (U, H) \rightarrow (\tilde{U}, \tilde{H})$ be a Heisenberg diffeomorphism. Let $\tilde{P} \in \Psi_{\tilde{H}}^m(V)$ and set $P = \phi^* \tilde{P}$. We need to show that P is a Ψ_H DO of order m on U .

First, by Proposition 3.16 the distribution kernel of $\tilde{P}$ takes the form,

$$(8.1) \quad k_{\tilde{P}}(\tilde{x}, \tilde{y}) = |\varepsilon'_x| K_{\tilde{P}}(\tilde{x}, -\varepsilon_{\tilde{x}}(\tilde{y})) + \tilde{R}(\tilde{x}, \tilde{y}),$$

with $K_{\tilde{P}}(\tilde{x}, \tilde{y})$ in $\mathcal{K}^{\hat{m}}(\tilde{U} \times \mathbb{R}^{d+1})$ and $\tilde{R}(\tilde{x}, \tilde{y})$ in $C^\infty(\tilde{U} \times \tilde{U})$. Therefore, the distribution kernel of $P = \phi^* \tilde{P}$ is given by

$$(8.2) \quad k_P(x, y) = |\phi'(y)| k_{\tilde{P}}(\phi(x), \phi(y)) = |\varepsilon'_x| K(x, -\varepsilon_x(y)) + \tilde{R}(\phi(x), \phi(y)),$$

where K is the distribution on $\mathcal{U} = \{(x, y) \in U \times \mathbb{R}^{d+1}; \varepsilon_x^{-1}(-y) \in U\} \subset U \times \mathbb{R}^{d+1}$ given by

$$(8.3) \quad K(x, y) = |\partial_y \Phi(x, y)| K_{\tilde{P}}(\phi(x), \Phi(x, y)), \quad \Phi(x, y) = -\varepsilon_{\phi(x)} \circ \phi \circ \varepsilon_x^{-1}(-y).$$

Next, it follows from [Po2, Props. 3.16, 3.18] that we have

$$(8.4) \quad \Phi(x, y) = -\phi'_H(x)(-y) + \Theta(x, y) = \phi'_H(x)(y) + \Theta(x, y),$$

where $\Theta(x, y)$ is a smooth function on $\mathcal{U}$ with a behavior near $y = 0$ of the form

$$(8.5) \quad \Theta(x, y) = (O(\|y\|^3), O(\|y\|^2), \dots, O(\|y\|^2)).$$

Then a Taylor expansion around $\tilde{y} = \phi'_H(x)y$ gives

$$(8.6) \quad K(x, y) = \sum_{\langle \alpha \rangle < N} |\partial_y \Phi(x, y)| \frac{\Theta(x, y)^\alpha}{\alpha!} (\partial_{\tilde{y}}^\alpha K_{\tilde{P}})(\phi(x), \phi'_H(x)y) + R_N(x, y),$$

$$(8.7) \quad R_N(x, y) = \sum_{\langle \alpha \rangle = N} |\partial_y \Phi(x, y)| \frac{\Theta(x, y)^\alpha}{\alpha!} \int_0^1 (t-1)^{N-1} \partial_{\tilde{y}}^\alpha K_{\tilde{P}}(\phi(x), \Phi_t(x, y)) dt.$$

where we have let $\Phi_t(x, y) = \phi'_H(x)y + t\Theta(x, y)$.

Set $f_\alpha(x, y) = |\partial_y \Phi(x, y)| \Theta(x, y)^\alpha$. Then (8.5) implies that near $y = 0$ we have

$$(8.8) \quad f_\alpha(x, y) = O(\|y\|^{3\alpha_0+2(\alpha_1+\dots+\alpha_d)}) = O(\|y\|^{\frac{3}{2}\langle \alpha \rangle}).$$

Thus all the homogeneous components of degree $< \frac{3}{2}\langle \alpha \rangle$ in the Taylor expansion for $f_\alpha(x, y)$ at $y = 0$ must be zero. Therefore, we can write

$$(8.9) \quad f_\alpha(x, y) = \sum_{\frac{3}{2}\langle \alpha \rangle \leq \langle \beta \rangle < \frac{3}{2}N} f_{\alpha\beta}(x) \frac{y^\beta}{\beta!} + \sum_{\langle \beta \rangle \doteq \frac{3}{2}N} r_{N\alpha\beta}(x, y) y^\beta,$$

where we have let $f_{\alpha\beta}(x) = \partial_y^\beta f_\alpha(x, 0)$, the functions $r_{M\alpha\beta}(x, y)$ are in $C^\infty(\mathcal{U})$ and the notation $\langle \beta \rangle \doteq \frac{3}{2}N$ means that $\langle \beta \rangle$ is equal to $\frac{3}{2}N$ if $\frac{3}{2}N$ is an integer and to $\frac{3}{2}N + \frac{1}{2}$ otherwise. Thus,

$$(8.10) \quad K(x, y) = \sum_{\langle \alpha \rangle < N} \sum_{\frac{3}{2}\langle \alpha \rangle \leq \langle \beta \rangle < \frac{3}{2}N} K_{\alpha\beta}(x, y) + \sum_{\langle \alpha \rangle < N} R_{N\alpha}(x, y) + R_N(x, y),$$

where we have let

$$(8.11) \quad K_{\alpha\beta}(x, y) = f_{\alpha\beta}(x) y^\beta (\partial_{\tilde{y}}^\alpha K_{\tilde{P}})(\phi(x), \phi'_H(x)y),$$

$$(8.12) \quad R_{N\alpha}(x, y) = \sum_{\langle \beta \rangle \doteq \frac{3}{2}N} r_{M\alpha\beta}(x, y) y^\beta (\partial_{\tilde{y}}^\alpha K_{\tilde{P}})(\phi(x), \phi'_H(x)y).$$

As in the proof of Proposition 3.16 the smoothness of $\phi'_H(x)y$ and the fact that $\phi'_H(x)(\lambda.y) = \lambda.\phi_H(x)y$ for any $\lambda \in \mathbb{R}$ imply that $K_{\alpha\beta}(x, y)$ belongs to $\mathcal{K}^{\hat{m}-\langle \alpha \rangle + \langle \beta \rangle}(U \times \mathbb{R}^{d+1})$. Notice that if $\frac{3}{2}\langle \alpha \rangle \leq \langle \beta \rangle \doteq \frac{3}{2}N$ then $\Re \hat{m} - \langle \alpha \rangle + \langle \beta \rangle \geq \Re \hat{m} + \frac{1}{3}\langle \beta \rangle \geq \Re \hat{m} + \frac{N}{2}$. It thus follows from Lemma 8.1 that, for any integer J , the remainder term $R_{N\alpha}$ is in $C^J(U \times \mathbb{R}^{d+1})$ as soon as N is large enough.

Let $\pi_x : U \times \mathbb{R}^{d+1} \rightarrow U$ denote the projection on the first coordinate. In the sequel we will say that a distribution $K(x, y) \in \mathcal{D}'(U \times \mathbb{R}^{d+1})$ is properly supported with respect to x when $\pi_x|_{\text{supp } K}$ is a proper map, i.e. for any compact $L \subset U$ the set $\text{supp } K \cap (L \times \mathbb{R}^{d+1})$ is compact.

In order to deal with the regularity of $R_N(x, y)$ in (8.7) we need the lemma below.

Lemma 8.2. *There exists a function $\chi_n \in C_c^\infty(\mathcal{U})$ properly supported with respect to x such that $\chi(x, y) = 1$ near $y = 0$ and, for any multi-order α , we can write*

$$(8.13) \quad \chi(x, y) \Theta(x, y)^\alpha = \sum_{\langle \beta \rangle \doteq \frac{3}{2}\langle \alpha \rangle} \theta_{\alpha\beta}(t, x, y) \Phi_t(x, y)^\beta$$

where the functions $\theta_{\alpha\beta}(t, x, y)$ are in $C^\infty([0, 1] \times U \times \mathbb{R}^{d+1})$.

Proof of the lemma. Let U' be a relatively compact open subset of U and let $(t_0, x_0) \in [0, 1] \times U'$. Since $\Phi_{t_0}(x_0, 0) = 0$ and $\partial_y \Phi_{t_0}(x_0, 0) = \phi'_H(x_0)$ is invertible the implicit function theorem implies that there exist an open interval I_{x_0} containing t_{x_0} , an open subset U_{x_0} of U containing x_0 , open subsets V_{x_0} and $\tilde{V}_{x_0}$ of $\mathbb{R}^{d+1}$ containing 0 and a smooth map $\Psi_{x_0}(t, x, \tilde{y})$ from $I_{x_0} \times U_{x_0} \times \tilde{V}_{x_0}$ to V_{x_0} such that $U_{x_0} \times V_{x_0}$ is contained in $\mathcal{U}$ and for any (t, x, y) in $I_{x_0} \times U_{x_0} \times V_{x_0}$ and any $\tilde{y}$ in $\tilde{V}_{x_0}$ we have

$$(8.14) \quad \tilde{y} = \Phi_t(x, y) \iff y = \Psi_{x_0}(t, x, \tilde{y}).$$

Since $[0, 1] \times \overline{U'}$ is compact we can cover it by finitely many products $I_{x_k} \times U_{x_k}$, $k = 1, \dots, p$, with $(t_k, x_k) \in [0, 1] \times \overline{U'}$. In particular, the sets $I = \cup_k I_k$ and $U'' = \cup U_k$ are open neighborhoods of I and $\overline{U'}$ respectively. Thanks to (8.14) we have $\Psi_{x_k} = \Psi_{x_l}$ on $(I_{x_k} \times U_{x_k} \times V_{x_k}) \cap (I_{x_l} \times U_{x_l} \times V_{x_l})$. Therefore, setting $V = \cap_k V_k$ and $\tilde{V} = \cap_k \tilde{V}_k$ we have $U'' \times V \subset \mathcal{U}$ and there exists a smooth map Ψ from $I \times U'' \times \tilde{V}$ such that for any (t, x, y) in $I \times U'' \times V$ and any $\tilde{y}$ in $\tilde{V}$ we have

$$(8.15) \quad \tilde{y} = \Phi_t(x, y) \iff y = \Psi(t, x, \tilde{y}).$$

Furthermore, as $\partial_{\tilde{y}} \Psi(t, x, 0) = [\partial_y \Phi_t(x, 0)]^{-1} = \phi'_H(x)^{-1}$ and $\phi'_H(x)^{-1}(\lambda y) = \lambda \phi'_H(x)^{-1}(y)$ for any $\lambda \in \mathbb{R}$, the function $\Theta(x, \Psi(t, x, \tilde{y}))$ behaves near $\tilde{y} = 0$ as in (8.5). Therefore, as in (8.8)–(8.9) for any multi-order α we can write

$$(8.16) \quad \Theta(x, \Psi(t, x, \tilde{y}))^\alpha = \sum_{\langle \beta \rangle \doteq \frac{3}{2} \langle \alpha \rangle} \tilde{\theta}_{\alpha\beta}(t, x, \tilde{y}) \tilde{y}^\beta, \quad \tilde{\theta}_{\alpha\beta}(t, x, \tilde{y}) \in C^\infty(I \times U'' \times \tilde{V}).$$

Setting $\tilde{y} = \Phi_t(x, y)$ then gives

$$(8.17) \quad \Theta(x, y)^\alpha = \sum_{\langle \beta \rangle \doteq \frac{3}{2} \langle \alpha \rangle} \theta_{\alpha\beta}(t, x, y) \Phi_t(x, y)^\beta, \quad \theta_{\alpha\beta}(t, x, y) \in C^\infty(I \times U'' \times V).$$

All this allows us to construct locally finite coverings $(U'_n)_{n \geq 0}$ and $(U''_n)_{n \geq 0}$ of U by relatively compact open subsets in such way that, for each integer n , the open U''_n contains $\overline{U'_n}$ and there exists an open $V_n \subset \mathbb{R}^{d+1}$ containing 0 so that, for any multiorder α , on $[0, 1] \times U''_n \times V_n$ we have

$$(8.18) \quad \Theta(x, y)^\alpha = \sum_{\langle \beta \rangle \doteq \frac{3}{2} \langle \alpha \rangle} \theta_{\alpha\beta}^{(n)}(t, x, y) \Phi_t(x, y)^\beta, \quad \theta_{\alpha\beta}^{(n)}(t, x, y) \in C^\infty([0, 1] \times U''_n \times V_n).$$

For each n let $\varphi_n \in C_c^\infty(U''_n)$ be such that $\varphi_n = 1$ on U'_n and let $\psi_n \in C_c^\infty(V_n)$ be such that $\psi_n = 1$ on a neighborhood V'_n of 0. Then we construct a locally finite family $(\chi_n)_{n \geq 0} \subset C_c^\infty(\mathcal{U})$ as follows: for $n = 0$ we set $\chi_0(x, y) = \varphi_0(x) \psi_0(y)$ and for $n \geq 1$ we let

$$(8.19) \quad \chi_n(x, y) = (1 - \varphi_0(x) \psi_0(y)) \dots (1 - \varphi_{n-1}(x) \psi_{n-1}(y)) \varphi_n(x) \psi_n(y).$$

Then $\chi = \sum \chi_n$ is a well defined smooth function on $U \times \mathbb{R}^{d+1}$ supported on $\cup_{n \geq 0} (U''_n \times V_n) \subset \mathcal{U}$, hence properly supported with respect to x . Also, as $\chi(x, y) = 1$ on each product $U'_n \times V'_n$ we see that $\chi(x, y) = 1$ on a neighborhood of $U \times \{0\}$. In addition, thanks to (8.18) on $[0, 1] \times \mathcal{U}$ we have

$$(8.20) \quad \chi(x, y) \Theta(x, y)^\alpha = \sum_{n \geq 0} \chi_n(x, y) \Theta(x, y)^\alpha = \sum_{\langle \beta \rangle \doteq \frac{3}{2} \langle \alpha \rangle} \theta_{\alpha\beta}(t, x, y) \Phi_t(x, y)^\beta,$$

where the functions $\theta_{\alpha\beta}(t, x, y) := \sum_n \chi_n(x, y) \theta_{\alpha\beta}^{(n)}(t, x, y)$ are in $C^\infty([0, 1] \times U \times \mathbb{R}^{d+1})$. The lemma is thus proved. $\square$

Let us go back to the proof of Proposition 3.18. Thanks to (8.7) and (8.13) on $U \times \mathbb{R}$ we have

$$(8.21) \quad \chi(x, y) R_N(x, y) = \sum_{\langle \alpha \rangle = N} \sum_{\langle \beta \rangle \doteq \frac{3}{2} N} \int_0^1 r_{\alpha\beta}(t, x, y) (\tilde{y}^\beta \partial_{\tilde{y}}^\alpha K_{\tilde{P}})(\phi(x), \Phi_t(x, y)) dt,$$

for some functions $r_{\alpha\beta}(t, x, y)$ in $C^\infty([0, 1] \times U \times \mathbb{R}^{d+1})$. Since $\tilde{y}^\beta \partial_{\tilde{y}}^\alpha K_{\tilde{P}}$ belongs to $\mathcal{K}^{\hat{m}+N/2}(\tilde{U} \times \mathbb{R}^{d+1})$ it follows from Lemma 8.1 that, for any integer $J \geq 0$, as soon as N is taken large enough $\tilde{y}^\beta \partial_{\tilde{y}}^\alpha K_{\tilde{P}}$ is in $C^J(\tilde{U} \times \mathbb{R}^{d+1})$ and so $\chi(x, y) R_N(x, y)$ is in $C^J(U \times \mathbb{R}^{d+1})$.

On the other hand, set $K_P(x, y) = \chi(x, y) K(x, y) = \sum \chi_n(x, y) K(x, y)$. Since $\chi(x, y)$ is supported in $\mathcal{U}$ and is properly supported with respect to x this defines a distribution on $U \times \mathbb{R}^{d+1}$.

Moreover, using (8.10) we get

$$(8.22) \quad K_P(x, y) = \sum_{\langle \alpha \rangle < N} \sum_{\frac{3}{2}\langle \alpha \rangle \leq \langle \beta \rangle < \frac{3}{2}N} K_{\alpha\beta}(x, y) + \sum_{j=1}^3 R_N^{(j)},$$

where the remainder terms $R_{N,z}^{(j)}$, $j = 1, 2, 3$ are given by

$$(8.23) \quad R_N^{(1)} = \chi(x, y)R_N(x, y), \quad R_N^{(2)} = \sum_{\langle \alpha \rangle < N} \chi(x, y)R_{N\alpha}(x, y),$$

$$(8.24) \quad R_N^{(2)}(x, y) = \sum_{\langle \alpha \rangle < N} \sum_{\frac{3}{2}\langle \alpha \rangle \leq \langle \beta \rangle < \frac{3}{2}N} (1 - \chi(x, y))K_{\alpha\beta}(x, y).$$

Each term $K_{\alpha\beta}(x, y)$ belongs to $\mathcal{K}^{\hat{m}-\langle \alpha \rangle + \langle \beta \rangle}(U \times \mathbb{R}^{d+1})$ and, as $\hat{m} + \langle \beta \rangle - \langle \alpha \rangle = \hat{m} + j$ and $\frac{3}{2}\langle \alpha \rangle \leq \langle \beta \rangle$ imply $\langle \alpha \rangle \leq 2j$ and $\langle \beta \rangle \leq \frac{4}{3}j$, in the r.h.s. (8.22) there are only finitely many such distributions in a given space $\mathcal{K}^{\hat{m}+j}(U \times \mathbb{R}^{d+1})$ as α and β range over all multi-orders such that $\frac{3}{2}\langle \alpha \rangle \leq \langle \beta \rangle$.

Furthermore, the reminder term $R_N^{(3)}$ is smooth and the other remainder terms $R_N^{(j)}$, $j = 1, 2$, are in $C^J(U \times \mathbb{R}^{d+1})$ as soon as N is large enough. Thus,

$$(8.25) \quad K_P(x, y) \sim \sum_{\frac{3}{2}\langle \alpha \rangle \leq \langle \beta \rangle} K_{\alpha\beta}(x, y),$$

which implies that K_P belongs to $\mathcal{K}^{\hat{m}}(U \times \mathbb{R}^{d+1})$ and satisfies (8.25).

Finally, from (8.1) and the very definition of $\Phi(x, y)$ on $U \times U$, we deduce that the distribution kernel of P is equal to

$$(8.26) \quad |\varepsilon'_x|K_P(x, -\varepsilon_x(y)) + [1 - \chi(x, \varepsilon_x(y))]| \varepsilon'_{\phi(x)}|K_{\tilde{P}}(\phi(x), -\tilde{\varepsilon}_{\phi(x)} \circ \phi(y)) + \tilde{R}(\phi(x), \phi(y)) \\ = |\varepsilon'_x|K_P(x, -\varepsilon_x(y)) \quad \text{mod } C^\infty(U \times U).$$

Combining this with Proposition 3.16 and the fact that $K_P(x, y)$ satisfies (8.25) completes the proof of Proposition 3.18.

9. PROOF OF PROPOSITION 3.21

Let $P : C_c^\infty(U) \rightarrow C^\infty(U)$ be a Ψ_H DO of order m and let us show that its transpose operator $P^t : C_c^\infty(U) \rightarrow C^\infty(U)$ is a Ψ_H DO of order m . By Proposition 3.16 the distribution kernel of P is of the form,

$$(9.1) \quad k_P(x, y) = |\varepsilon'_x|K_P(x, -\varepsilon_x(y)) + R(x, y),$$

with $K_P(x, y)$ in $\mathcal{K}^m(U \times \mathbb{R}^{d+1})$ and $R(x, y)$ in $C^\infty(U \times U)$. Thus the distribution kernel of P^t can be written as

$$(9.2) \quad k_{P^t}(x, y) = k_P(y, x) = |\varepsilon'_y|K_P(y, -\varepsilon_y(x)) + R(y, x) = |\varepsilon'_x|K(x, -\varepsilon_x(y)) + R(y, x),$$

where K is the distribution on the open $\mathcal{U} = \{(x, y); \varepsilon_x^{-1}(-y) \in U\}$ given by

$$(9.3) \quad K(x, y) = |\varepsilon'_x|^{-1}|\varepsilon'_y|K_P(\varepsilon_x^{-1}(-y), -\varepsilon_{\varepsilon_x^{-1}(-y)}(x)).$$

Lemma 9.1. *On $\mathcal{U}$ we have*

$$(9.4) \quad \varepsilon_{\varepsilon_x^{-1}(-y)}(x) = y - \Theta(x, y),$$

where $\Theta : \mathcal{U} \rightarrow \mathbb{R}^{d+1}$ is a smooth map with a behavior near $y = 0$ of the form (8.5).

Proof. Let $(x, y) \in \mathcal{U}$ and $Y \in G_x U$ let $\lambda_y(Y) = y.Y$, that is, λ_y is the left multiplication by y on $G_x U$. Then by [Po2, Eq. (3.32)] for Y small enough we have

$$(9.5) \quad \lim_{t \rightarrow 0} \varepsilon_x \circ \varepsilon_x^{-1}(t, -y)(t.Y) = \lambda_{-y}(Y) = \lambda_y^{-1}(Y).$$

Since $\varepsilon_x \circ \varepsilon_x^{-1}(t, -y)(t.Y)$ is a smooth function of (t, Y) near $(0, 0)$, it follows from the implicit function theorem that for Y small enough we have

$$(9.6) \quad \lim_{t \rightarrow 0} t^{-1} \cdot \varepsilon_x^{-1}(t, -y) \circ \varepsilon_x^{-1}(Y) = \lambda_y(Y).$$

In particular, for $Y = 0$ we get

$$(9.7) \quad \lim_{t \rightarrow 0} t^{-1} \cdot \varepsilon_x^{-1}(t, -y)(x) = y.$$

Now, the function $\varepsilon_x^{-1}(-y)(x)$ depends smoothly on $(x, y) \in \mathcal{U}$, so (9.7) allows us to put it into the form,

$$(9.8) \quad \varepsilon_x^{-1}(-y)(x) = y - \Theta(x, y),$$

where $\Theta = (\Theta_0, \dots, \Theta_d)$ is smooth map from $\mathcal{U}$ to $\mathbb{R}^{d+1}$ with a behavior near $y = 0$ of the form

$$(9.9) \quad \Theta_0(x, y) = O(|y_0|^2 + |y_0||y| + |y|^3), \quad \Theta_j(x, y) = O(|y|^2), \quad j = 1, \dots, d.$$

In particular, near $y = 0$ the map Θ has a behavior of the form (8.5). $\square$

Next, a Taylor expansion around $(\varepsilon_x^{-1}(-y), -y)$ gives

$$(9.10) \quad K(x, y) = \sum_{\langle \alpha \rangle < N} |\varepsilon'_x|^{-1} |\varepsilon'_y| \frac{\theta(x, y)^\alpha}{\alpha!} (\partial_y^\alpha K_P)(\varepsilon_x^{-1}(-y), -y) + R_N(x, y),$$

$$(9.11) \quad R_N(x, y) = \sum_{\langle \alpha \rangle = N} |\varepsilon'_x|^{-1} |\varepsilon'_y| \frac{\theta(x, y)^\alpha}{\alpha!} \int_0^1 (1-t)^{N-1} \partial_y^\alpha K_P(\varepsilon_x^{-1}(-y), \Phi_t(x, y)),$$

where we have let $\Phi_t(x, y) = -y + t\Theta(x, y)$.

Let $a_\alpha(x, y) = |\varepsilon'_x|^{-1} |\varepsilon'_y| \frac{\theta(x, y)^\alpha}{\alpha!}$. Then because of (8.5) the same arguments used to prove (8.9) show that there exist functions $r_{N\alpha}(x, y) \in C^\infty(\mathcal{U})$, $\langle \beta \rangle \leq \frac{3}{2}N$, such that

$$(9.12) \quad a_\alpha(x, y) = \sum_{\frac{3}{2}\langle \alpha \rangle \leq \langle \beta \rangle < \frac{3}{2}N} a_{\alpha\beta}(x) y^\beta + \sum_{\langle \beta \rangle \geq \frac{3}{2}N} r_{N\alpha}(x, y) y^\beta,$$

where we have let $a_{\alpha\beta}(x) = \frac{1}{\beta!} \partial^\beta a_\alpha(x, 0)$. Thus,

$$(9.13) \quad K(x, y) = \sum_{\langle \alpha \rangle < N} \sum_{\frac{3}{2}\langle \alpha \rangle \leq \langle \beta \rangle < \frac{3}{2}N} f_{\alpha\beta}(x) y^\beta (\partial_y^\alpha K_P)(\varepsilon_x^{-1}(-y), -y) + \sum_{\langle \alpha \rangle < N} R_{N\alpha}(x, y) + R_N(x, y),$$

$$(9.14) \quad R_{N\alpha}(x, y) = \sum_{\langle \beta \rangle \geq \frac{3}{2}N} r_{N\alpha}(x, y) y^\beta (\partial_y^\alpha K_P)(\varepsilon_x^{-1}(-y), -y).$$

Next, a further Taylor expansion shows that $(\partial_y^\alpha K_P)(\varepsilon_x^{-1}(-y), -y)$ is equal to

$$(9.15) \quad \sum_{|\gamma| < N} \frac{1}{\gamma!} (\varepsilon_x^{-1}(-y) - x)^\gamma (\partial_x^\gamma \partial_y^\alpha K_P)(x, -y) + \sum_{|\gamma| = N} \int_0^1 (1-t)^{N-1} (\partial_x^\gamma \partial_y^\alpha K_P)(\varepsilon_t(x, y), -y),$$

where we have let $\varepsilon_t(x, y) = x + t(\varepsilon_x^{-1}(-y) - x)$. Since $\varepsilon_x^{-1}(-y) - x$ is polynomial in y of degree 2 and vanishes for $y = 0$, we can write

$$(9.16) \quad \frac{1}{\gamma!} (\varepsilon_x^{-1}(-y) - x)^\gamma = \sum_{|\gamma| \leq |\delta| \leq 2|\gamma|} a_{\gamma\delta}(x) y^\delta, \quad b_{\gamma\delta}(x) = \frac{1}{\gamma! \delta!} [\partial_y (\varepsilon_x^{-1}(-y) - x)^\gamma]_{y=0}.$$

Therefore, we can put $K(x, y)$ into the form,

$$(9.17) \quad K(x, y) = \sum_{\alpha, \beta, \gamma, \delta}^{(N)} K_{\alpha\beta\gamma\delta}(x, y) + \sum_{\langle\alpha\rangle < N} \sum_{\frac{3}{2}\langle\alpha\rangle \leq \langle\beta\rangle < \frac{3}{2}N} R_{N\alpha\beta}(x, y) + \sum_{\langle\alpha\rangle < N} R_{N\alpha}(x, y) + R_N(x, y),$$

where the first summation goes over all the multi-orders α, β, γ and δ such that $\langle\alpha\rangle < N, \frac{3}{2}\langle\alpha\rangle \leq \langle\beta\rangle < \frac{3}{2}N$ and $|\gamma| \leq |\delta| \leq 2|\gamma| < 2N$, and we have let

$$(9.18) \quad K_{\alpha\beta\gamma\delta}(x, y) = a_{\alpha\beta\gamma\delta}(x) y^{\beta+\delta} (\partial_x^\gamma \partial_y^\alpha K_P)(x, -y), \quad f_{\alpha\beta\gamma\delta}(x) = a_{\alpha\beta}(x) b_{\gamma\delta}(x),$$

$$(9.19) \quad R_{N\alpha\beta}(x, y) = \sum_{|\gamma|=N} \sum_{N \leq |\delta| \leq 2N} a_{\alpha\beta\gamma\delta}(x) y^{\beta+\delta} \int_0^1 (1-t)^{N-1} (\partial_x^\gamma \partial_y^\alpha K_P)(\varepsilon_t(x, y), -y).$$

Now, the distribution $y^\beta K_P(x, -y)$ belongs to $\mathcal{K}^{\hat{m}-\langle\alpha\rangle+\langle\beta\rangle}(U \times \mathbb{R}^{d+1})$. In particular, if $\frac{3}{2}\langle\alpha\rangle \leq \langle\beta\rangle \leq \frac{3}{2}N$ then $\Re \hat{m} - \langle\alpha\rangle + \langle\beta\rangle \geq \Re \hat{m} + \frac{1}{3}\langle\beta\rangle \geq \Re \hat{m} + \frac{1}{2}N$. Therefore, for any given integer J Lemma 8.1 tells us that $y^\beta K_P(x, -y)$ is in $C^J(U \times \mathbb{R}^{d+1})$ as soon as N is large enough. It follows that all the remainder terms $R_{N\alpha}(x, y), \langle\alpha\rangle < N$, belong to $C^J(\mathcal{U})$ for N large enough.

Similarly, if $\frac{3}{2}\langle\alpha\rangle \leq \langle\beta\rangle$ and $|\gamma| = N \leq |\delta| \leq 2N$ then $\Re \hat{m} - \langle\alpha\rangle + \langle\beta\rangle + \langle\delta\rangle \geq \Re \hat{m} + \langle\delta\rangle \geq \Re \hat{m} + \frac{1}{2}N$, so using Lemma 8.1 we see that $y^{\beta+\delta} (\partial_x^\gamma \partial_y^\alpha K_P)(x, -y)$ is in $C^J(U \times \mathbb{R}^{d+1})$ for N large enough. It then follows that for N large enough the remainder terms $R_{N\alpha\beta}(x, y)$ with $\langle\alpha\rangle < N$ and $\frac{3}{2}\langle\alpha\rangle \leq \langle\beta\rangle \leq \frac{3}{2}N$ are all in $C^J(\mathcal{U})$ as soon as N is chosen large enough.

In order to deal with the last remainder term $R_N(x, y)$ notice that, along the same lines as that of the proof of Lemma 8.2, one can show that there exists a $\chi \in C^\infty(\mathcal{U})$ properly supported with respect to x such that $\chi(x, y) = 1$ near $y = 0$ and, for any multi-order α , we can write

$$(9.20) \quad \chi(x, y) \Theta(x, y)^\alpha = \sum_{\langle\beta\rangle \doteq \frac{3}{2}\langle\alpha\rangle} \theta_{\alpha\beta}(t, x, y) \Phi_t(x, y)^\beta,$$

where the functions $\theta_{\alpha\beta}(t, x, y)$ are in $C^\infty([0, 1] \times U \times \mathbb{R}^{d+1})$. Then we can put $\chi(x, y) R_N(x, y)$ into the form,

$$(9.21) \quad \chi(x, y) R_N(x, y) = \sum_{\langle\alpha\rangle = N} \sum_{\langle\beta\rangle \doteq \frac{3}{2}\langle\alpha\rangle} |\varepsilon'_x|^{-1} |\varepsilon'_y| \int_0^1 r_{N\alpha\beta}(t, x, y) (y^\beta \partial_y^\alpha K_P)(\varepsilon_x^{-1}(-y), \Phi_t(x, y)),$$

for some functions $r_{N\alpha\beta}(t, x, y)$ in $C^\infty([0, 1] \times U \times \mathbb{R}^{d+1})$. As $(y^\beta \partial_y^\alpha K_P)$ is in $\mathcal{K}^{\hat{m}-\langle\alpha\rangle+\langle\beta\rangle}(U \times \mathbb{R}^{d+1})$ and we have $\Re \hat{m} - \langle\alpha\rangle + \langle\beta\rangle \geq \Re \hat{m} + \frac{1}{2}\langle\alpha\rangle = \Re \hat{m} + \frac{1}{2}N$, using Lemma 8.1 we see that for N large enough $\chi(x, y) R_N(x, y)$ is in $C^J(\mathcal{U})$, so is in $C^J(U \times \mathbb{R}^{d+1})$ since $\chi(x, y) R_N(x, y)$ is a properly supported with respect to x .

Let $K_{Pt}(x, y) = \chi(x, y) K(x, y)$. This defines a distribution on $U \times \mathbb{R}^{d+1}$ since χ is properly supported with respect to x . Moreover, we have

$$(9.22) \quad K_{Pt}(x, y) = \sum_{\alpha, \beta, \gamma, \delta}^{(N)} K_{\alpha\beta\gamma\delta}(x, y) + \sum_{j=1}^4 R_N^{(j)}(x, y),$$

where the remainder terms $R_N^{(j)}, j = 1, \dots, 4$, are given by

$$(9.23) \quad R_N^{(1)} = \chi(x, y) R_N(x, y), \quad R_N^{(2)} = \sum_{\langle\alpha\rangle < N} \chi(x, y) R_{N\alpha}(x, y),$$

$$(9.24) \quad R_N^{(3)} = \sum_{\langle\alpha\rangle < N} \sum_{\frac{3}{2}\langle\alpha\rangle \leq \langle\beta\rangle < \frac{3}{2}N} \chi(x, y) R_{N\alpha\beta}(x, y), \quad R_N^{(4)} = \sum_{\alpha, \beta, \gamma, \delta}^{(N)} (1 - \chi(x, y)) K_{\alpha\beta\gamma\delta}(x, y).$$

Note that $K_{\alpha\beta\gamma\delta}(x, y)$ belongs to $\mathcal{K}^{\hat{m}-\langle\alpha\rangle+\langle\gamma\rangle+\langle\delta\rangle}(U \times \mathbb{R}^{d+1})$ and there are finitely many terms of a given order as α, β, γ and δ ranges over all the multi-orders such that $\frac{3}{2}\langle\alpha\rangle \leq \langle\beta\rangle$ and $|\gamma| \leq |\delta| \leq 2|\gamma|$.

On the other hand, the remainder term $R_N^{(4)}$ is smooth and it follows from the observations above that the other remainder terms are in $C^J(U \times \mathbb{R}^{d+1})$ as soon as N is large enough. Thus,

$$(9.25) \quad K_{P^t}(x, y) \sim \sum_{\frac{3}{2}\langle\alpha\rangle \leq \langle\beta\rangle} \sum_{|\gamma| \leq |\delta| \leq 2|\gamma|} K_{\alpha\beta\gamma\delta}(x, y),$$

which incidentally shows that $K_{P^t}(x, y)$ belongs to $\mathcal{K}^{\hat{m}}(U \times \mathbb{R}^{d+1})$.

Finally, thanks to (9.2) we can put the kernel of P^t into the form,

$$(9.26) \quad k_{P^t}(x, y) = |\varepsilon'_x| K_P(x, -\varepsilon_x(y)) + |\varepsilon'_x| [(1 - \chi)K](x, -\varepsilon_x(y)) + R(y, x) \\ = |\varepsilon'_x| K_P(x, -\varepsilon_x(y)) \mod C^\infty(U \times U).$$

It then follows from Proposition 3.16 that P^t is a Ψ_H DO of order m . Moreover, working out the expression for $K_{\alpha\beta\gamma\delta}$ shows that the asymptotics expansion (9.25) reduces to (3.37). The proof of Proposition 3.21 is thus achieved.

REFERENCES

- [AS1] Atiyah, M.F.; Singer, I.M.: *The index of elliptic operators. I.* Ann. of Math. (2) **87** (1968), 484–530.
- [AS2] Atiyah, M.F.; Singer, I.M.: *The index of elliptic operators. III.* Ann. of Math. (2) **87**, 546–604 (1968).
- [APS] Atiyah, M., Patodi, V., Singer, I.: *Spectral asymmetry and Riemannian geometry. III.* Math. Proc. Camb. Philos. Soc. **79**, 71–99 (1976).
- [Be] Bellaïche, A.: *The tangent space in sub-Riemannian geometry.* Sub-Riemannian geometry, 1–78, Progr. Math., 144, Birkhäuser, Basel, 1996.
- [Bea] Beals, R.: *Opérateurs invariants hypoelliptiques sur un groupe de Lie nilpotent.* Séminaire Goulaouic-Schwartz (1976–77), Exposé No. 19. Ecole Polytech., Palaiseau, 1977.
- [BG] Beals, R.; Greiner, P.C.: *Calculus on Heisenberg manifolds.* Annals of Mathematics Studies, vol. 119. Princeton University Press, Princeton, NJ, 1988.
- [Bo1] Boutet de Monvel, L.: *Hypoelliptic operators with double characteristics and related pseudo-differential operators.* Comm. Pure Appl. Math. **27** (1974), 585–639.
- [Bo2] Boutet de Monvel, L.: *Logarithmic trace of Toeplitz projectors.* E-print, arXiv, Dec. 04.
- [BGH] Boutet de Monvel, L.; Grigis, A.; Helffer, B.: *Paramétrixes d'opérateurs pseudo-différentiels à caractéristiques multiples.* Journées EDP (Rennes, 1975) pp. 93–121. Astérisque, No. 34-35, Soc. Math. France, Paris, 1976.
- [BL] Brüning, J.; Lesch, M.: *On the eta-invariant of certain nonlocal boundary value problems.* Duke Math. J. **96** (1999), no. 2, 425–468.
- [CGGP] Christ, M.; Geller, D.; Głowacki, P.; Polin, L.: *Pseudodifferential operators on groups with dilations.* Duke Math. J. **68** (1992) 31–65.
- [Co] Connes, A.: *Noncommutative geometry.* Academic Press, Inc., San Diego, CA, 1994.
- [CM] Connes, A.; Moscovici, H.: *The local index formula in noncommutative geometry.* Geom. Funct. Anal. **5** (1995), no. 2, 174–243.
- [Dy1] Dynin, A.: *Pseudodifferential operators on the Heisenberg group.* Dokl. Akad. Nauk SSSR **225** (1975) 1245–1248.
- [Dy2] Dynin, A.: *An algebra of pseudodifferential operators on the Heisenberg groups. Symbolic calculus.* Dokl. Akad. Nauk SSSR **227** (1976), 792–795.
- [EMM] Epstein, C.L.; Melrose, R.B.; Mendoza, G.: *The Heisenberg algebra, index theory and homology.* To appear. Partially available at <http://www-math.mit.edu/~rbm>.
- [ET] Eliashberg, Y.; Thurston, W.: *Confoliations.* University Lecture Series, 13, AMS, Providence, RI, 1998.
- [Fe] Fefferman, C.: *Parabolic invariant theory in complex analysis.* Adv. in Math. **31** (1979), no. 2, 131–262.
- [Fo] Folland, G.: *Subelliptic estimates and function spaces on nilpotent Lie groups.* Ark. Mat. **13** (1975) 161–207.
- [FS] Folland, G.; Stein, E.: *Estimates for the $\bar{\partial}_b$ complex and analysis on the Heisenberg group.* Comm. Pure Appl. Math. **27** (1974) 429–522.
- [Ge] Liouville's theorem for homogeneous groups. Comm. Partial Differential Equations **8** (1983) 1665–1677.
- [Gi] P.B. Gilkey *Invariance theory, the heat equation, and the Atiyah-Singer index theorem.* Mathematics Lecture Series, 11. Publish or Perish, Inc., Wilmington, Del., 1984.

- [Gl] Głowacki, P.: *Stable semigroups of measures as commutative approximate identities on nongraded homogeneous groups*. Invent. Math. **83** (1986), no. 3, 557–582.
- [Gr] Greenleaf, A.: *The first eigenvalue of a sub-Laplacian on a pseudo-Hermitian manifold*. Comm. Partial Differential Equations **10** (1985), no. 2, 191–217.
- [Gri] Grigis, A.: *Hypoellipticité et paramétrix pour des opérateurs pseudodifférentiels à caractéristiques doubles*. Journées EDP (Rennes, 1975) pp. 183–205. Astérisque, No. 34-35, Soc. Math. France, Paris, 1976.
- [Gro] Gromov, M.: *Carnot-Carathéodory spaces seen from within. Sub-Riemannian geometry*, 79–323, Progr. Math., 144, Birkhäuser, Basel, 1996.
- [Gu] Guillemin, V.W.: *A new proof of Weyl’s formula on the asymptotic distribution of eigenvalues*. Adv. in Math. **55** (1985), no. 2, 131–160.
- [He] Helffer, B.: *Hypoellipticité pour des opérateurs différentiels sur des groupes de Lie nilpotents*. Pseudodifferential operator with applications (Bressanone, 1977), pp. 73–88, Liguori, Naples, 1978.
- [HN1] Helffer, B.; Nourrigat, F.: *Hypoellipticité pour des groupes nilpotents de rang de nilpotence 3*. Comm. Partial Differential Equations **3** (1978), no. 8, 643–743.
- [HN2] Helffer, B.; Nourrigat, J.: *Caractérisation des opérateurs hypoelliptiques homogènes invariants à gauche sur un groupe de Lie nilpotent gradué*. Comm. Partial Differential Equations **4** (1979), no. 8, 899–958.
- [HN3] Helffer, B.; Nourrigat, J.: *Hypoellipticité maximale pour des opérateurs polynômes de champs de vecteurs*. Prog. Math., No. 58, Birkhäuser, Boston, 1986.
- [Hi] Hirachi, K.: *Logarithmic singularity of the Szegő kernel and a global invariant of strictly pseudoconvex domains*. E-print, arXiv, Sep. 03, 17 pages. To appear in Ann. of Math..
- [Hö2] L. Hörmander *Hypoelliptic second order differential equations*. Acta Math. **119** (1967), 147–171.
- [Hö3] Hörmander, L.: *A class of hypoelliptic pseudodifferential operators with double characteristics*. Math. Ann. **217** (1975), no. 2, 165–188.
- [Ko] Koenig, K.: *On maximal Sobolev and Hlder estimates for the tangential Cauchy-Riemann operator and boundary Laplacian*. Amer. J. Math. **124** (2002), no. 1, 129–197.
- [Koh1] J. J. Kohn *Boundaries of complex manifolds*. 1965 Proc. Conf. Complex Analysis (Minneapolis, 1964) pp. 81–94 Springer, Berlin
- [Koh2] Kohn, J. J.: *Superlogarithmic estimates on pseudoconvex domains and CR manifolds*. Ann. of Math. (2) **156** (2002), no. 1, 213–248.
- [KR] Kohn, J.J.; Rossi, H. *On the extension of holomorphic functions from the boundary of a complex manifold*. Ann. of Math. **81** (1965) 451–472.
- [Le] Lee, J.M.: *The Fefferman metric and pseudo-Hermitian invariants*. Trans. Amer. Math. Soc. **296** (1986), no. 1, 411–429.
- [Me2] Melrose, R.B.: *The Atiyah-Patodi-Singer index theorem*. A.K. Peters, Boston, 1993.
- [Ni] Nicoara, A.: *Regularity results for the tangential Cauchy-Riemann operator and the Szegő projection on weakly pseudoconvex CR manifolds*. PhD dissertation, Princeton University, 2002.
- [Po1] Ponge, R.: *Spectral asymmetry, zeta functions and the noncommutative residue*. E-print, arXiv, Oct. 03, 28 pages. Submitted to J. Funct. Anal..
- [Po2] Ponge, R.: *The tangent groupoid of a Heisenberg manifold*. E-print, arXiv, Apr. 04, 17 pages.
- [Po3] Ponge, R.: *Functional calculus and spectral asymptotics for hypoelliptic operators on Heisenberg manifolds. I*. E-print, arXiv, Feb. 05, 74 pages.
- [Po4] Ponge, R.: *Functional calculus and spectral asymptotics for hypoelliptic operators on Heisenberg manifolds. II*. In preparation.
- [Ro1] Rockland, C.: *Hypoellipticity on the Heisenberg group-representation-theoretic criteria*. Trans. Amer. Math. Soc. **240** (1978) 1–52.
- [Ro2] Rockland, C.: *Intrinsic nilpotent approximation*. Acta Appl. Math. **8** (1987), no. 3, 213–270.
- [RSt] Rothschild, L.; Stein, E.: *Hypoelliptic differential operators and nilpotent groups*. Acta Math. **137** (1976) 247–320.
- [Ru] Rumin, M.: *Formes différentielles sur les variétés de contact*. J. Differential Geom. **39** (1994), no.2, 281–330.
- [Ta] Tanaka, N.: *A differential geometric study on strongly pseudo-convex manifolds*. Lectures in Mathematics, Department of Mathematics, Kyoto University, No. 9. Kinokuniya Book-Store Co., Ltd., Tokyo, 1975.
- [Tay] Taylor, M.E.: *Noncommutative microlocal analysis. I*. Mem. Amer. Math. Soc. **52** (1984), no. 313,
- [Wol] Wodzicki, M.: *Local invariants of spectral asymmetry*. Invent. Math. **75** (1984) 143–177.

DEPARTMENT OF MATHEMATICS, OHIO STATE UNIVERSITY, COLUMBUS, USA.
E-mail address: `ponge@math.ohio-state.edu`